

For Research Use Only

Phospho-MLKL (Ser358) Recombinant monoclonal antibody

Catalog Number: 82090-2-RR

2 Publications



Basic Information

Catalog Number:

82090-2-RR

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

XM_005255834

GeneID (NCBI):

197259

UNIPROT ID:

Q8NB16

Full Name:

mixed lineage kinase domain-like

Calculated MW:

54 kDa

Observed MW:

54 kDa

Purification Method:

Protein A purification

CloneNo.:

241469D2

Recommended Dilutions:

WB: 1:5000-1:50000

Applications

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity:

human

Cited Species:

mouse

Positive Controls:

WB : TNF alpha and Calyculin A treated HT-29 cells,

Background Information

Mixed lineage kinase domain like pseudokinase (MLKL), belongs to the protein kinase superfamily. One common method for inducing necroptosis is by treating cells grown in culture with a mixture composed of TNF- α , a Smac-mimetic, that degrades cellular inhibitors of apoptosis (cIAPs) and the pancaspase inhibitor Z-VAD-FMK. This mixture is abbreviated as T/S/Z. It drives necrosome complex formation and RIPK3-dependent phosphorylation of Thr-357 and Ser-358 located within the C-terminal kinase-like domain of human MLKL. (PMID: 28878015)

Notable Publications

Author	Pubmed ID	Journal	Application
Xuemei Yao	40344881	Biomaterials	WB
Kai Ma	40430756	Pathogens	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated HT-29 cells, TNF alpha and Calyculin A treated HT-29 cells were subjected to SDS PAGE followed by western blot with 82090-2-RR (Phospho-MLKL (Ser358) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Beta Actin (66009-1-Ig) antibody as a loading control.