For Research Use Only

IDH2 Recombinant antibody

Catalog Number:82117-1-RR

Featured Product



Basic Information

Catalog Number: GenBank Accession Number: 82117-1-RR BC009244

 Size:
 GeneID (NCBI):
 CloneNo.:

 1000 ug/ml
 3418
 1B1

 Source:
 UNIPROT ID:
 Recomme

 Source:
 UNIPROT ID:
 Recommended Dilutions:

 Rabbit
 P48735
 WB 1:5000-1:50000

 Isotype:
 Full Name:
 IP 0.5-4.0 ug for 1.0-3.0 mg of total

Isotype: Full Name: IP 0.5-4.0 ug for 1.

IgG isocitrate dehydrogenase 2 (NADP+), protein lysate
IHC 1:500-1:2000

Immunogen Catalog Number: mitochondrial

AG8779 Calculated MW: 452 aa, 51 kDa

Observed MW: 43 kDa

Applications

Tested Applications: WB, IHC, IP, ELISA

Species Specificity: human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: LNCaP cells, HEK-293 cells, mouse heart tissue,

Purification Method:

Protein A purification

Jurkat cells, rat heart tissue

IP: mouse brain tissue.

IHC: human colon cancer tissue,

Background Information

IDH2, also named as IDP and ICD-M, belongs to the isocitrate and isopropylmalate dehydrogenases family. It plays a role in intermediary metabolism and energy production. IDH2 is a mitochondrial NADP-dependent isocitrate dehydrogenase that catalyzes oxidative decarboxylation of isocitrate to alpha-ketoglutarate, producing NADPH. It may tightly associate or interact with the pyruvate dehydrogenase complex. IDH1 and IDH2 mutations could also contribute to tumorigenesis and cancer progression through increased mutagenesis. IDH2 has 2 isoforms with the MW of 51 kDa and 45 kDa, and the mature form is about 47 kDa and 41 kDa with the N-terminal transit peptide cleaved.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82117-1-RR (IDH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 82117-1-RR (IDH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of IDH2 antibody (82117-1-RR; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-IDH2 transfected HEK-293 cells.



IP result of anti-IDH2 (IP:82117-1-RR, 4ug; Detection:82117-1-RR 1:1000) with mouse brain tissue lysate 1280 ug.



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 82117-1-RR (IDH2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).