For Research Use Only

SIRT3 Recombinant antibody

Catalog Number:82210-1-RR





Purification Method: Protein A purification CloneNo.:			
		2P6 Recommended Dilutions: WB 1:5000-1:50000 IHC 1:200-1:800	
Positive Controls: WB : HeLa cells, mouse heart tissue, rat heart tissue, HepG2 cells, HT-29 cells, mouse liver tissue, pig live tissue, rat liver tissue IHC : human prostate cancer tissue,			
			T3 is the only sirtuin whose increased ns. It is localized in the mitochondrial uding acetyl coenzyme A synthetase 2 om genotoxic and oxidative stress- it is activated inside the mitochondria essed form of SIRT3 is approximately 2 nd is processed by mitochondrial

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data






Various lysates were subjected to SDS PAGE followed by western blot with 82210-1-RR (SIRT3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffinembedded human prostate cancer tissue slide using 82210-1-RR (SIRT3 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 82210-1-RR (SIRT3 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLL) kinetic assays of 82210-1-RR against Human SIRT3 were performed. The affinity constant is 0.261 nM.