## For Research Use Only

## ATP5A1 Recombinant antibody, PBS Only

Catalog Number:82288-1-PBS



**Purification Method:** 

Protein A purification

CloneNo.:

4012

**Basic Information** 

Catalog Number: GenBank Accession Number: 82288-1-PBS BC064562

 Size:
 GeneID (NCBI):

 1mg/ml
 498

Source: UNIPROT ID:
Rabbit P25705
Isotype: Full Name:

IgG ATP synthase, H+ transporting, mitochondrial F1 complex, alpha

AG6385 subunit 1, cardiac muscle

Calculated MW: 60 kDa

> Observed MW: 50-55 kDa

**Applications** 

**Tested Applications:** 

WB, IHC, IF/ICC, FC (Intra), Indirect ELISA

Species Specificity: human, mouse, rat

## **Background Information**

The ATP5A1 gene encodes the  $\,^{\alpha}$  subunit of mitochondrial ATP synthase which produces ATP from ADP in the presence of a proton gradient across the membrane. The mitochondrial ATP synthase, also known as Complex V or F1F0 ATP synthase, is a multi-subunit enzyme complex consisting of two functional domains, the F1-containing the catalytic core and the F0-containing the membrane proton channel. F0 domain has 10 subunits: a,b, c, d, e, f, g, OSCP, A6L, and F6. F1 is composed of subunits  $\,^{\alpha}$ ,  $\,^{\beta}$ ,  $\,^{\beta}$ ,  $\,^{\epsilon}$ , and a loosely attached inhibitor protein IF1. Recently defect in ATP5A1 has been linked to the fatal neonatal mitochondrial encephalopathy. ATP5A1 is localized in the mitochondria and anti-ATP5A1 can be used as the loading control for mitochondrial or Complex V proteins. This antibody recognizes the endogenous ATP5A1 protein in lysates from various cell lines and tissues. The predicted MW of ATP5A1 is 60 kDa, while it undergoes the transit peptide cleavage to become a mature form around 50-55 kDa. Several isoforms of ATP5A1 exist due to the alternative splicing.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

PBS Only

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82288-1-RR (ATP5A1 antibody) at dilution of 1:30000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82288-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ATP5A1 antibody (82288-1-RR, Clone: 4012) at dilution of 1:2000 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red). This data was developed using the same antibody clone with 82288-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82288-1-RR (ATP5A1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82288-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 82288-1-RR (ATP5A1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82288-1-PBS in a different storage buffer formulation.



1X10^6 HeLa cells were intracellularly stained with 0.4 ug Anti-Human ATP5A1 (82288-1-RR, Clone:4012) and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Isotype Control. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 82288-1-PBS in a different storage buffer



Biolayer interferometry (BLL) kinetic assays of 82288-1-RR against Human ATP5A1 were performed. The affinity constant is 2.37 nM.