

For Research Use Only

Phospho-IkB Alpha (Ser32/36) Recombinant antibody

Catalog Number: 82349-1-RR

41 Publications



Basic Information

Catalog Number:

82349-1-RR

Concentration:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC004983

GeneID (NCBI):

4792

UNIPROT ID:

P25963

Full Name:

nuclear factor of kappa light
polypeptide gene enhancer in B-cells
inhibitor, alpha

Calculated MW:

36 kDa

Observed MW:

36 kDa

Purification Method:

Protein A purification

CloneNo.:

1G21

Recommended Dilutions:

WB 1:1000-1:4000

IHC 1:1000-1:4000

Applications

Tested Applications:

WB, IHC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse, rat, pig

**Note-IHC: suggested antigen retrieval with
TE buffer pH 9.0; (*) Alternatively, antigen
retrieval may be performed with citrate
buffer pH 6.0**

Positive Controls:

WB : Calyculin A treated HeLa cells, Calyculin A
treated HSC-T6 cells, Calyculin A treated NIH/3T3
cells, rat liver tissue, mouse liver tissue

IHC : human skin cancer tissue,

Background Information

Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha (NFKB1A, synonyms: IKBA, MAD-3, NFKB1). NFKB1 or NFKB2 is bound to REL, RELA, or RELB to form the NFKB complex. The NFKB complex is inhibited by I-kappa-B proteins (NFKBIA or NFKBIB), which inactivate NF-kappa-B by trapping it in the cytoplasm. Phosphorylation of serine residues on the I-kappa-B proteins by kinases (IKBKA, or IKBKB) marks them for destruction via the ubiquitination pathway, thereby allowing activation of the NF-kappa-B complex. Activated NFKB complex translocates into the nucleus and binds DNA at kappa-B-binding motifs such as 5-prime GGGRNYYCC 3-prime or 5-prime HGGARNYYCC 3-prime. The antibody also detects the phosphorylated form of the NFKBIA (IkB Alpha) (PMID: 16904979, PMID: 28990531 PMID: 10455908).

Notable Publications

Author	Pubmed ID	Journal	Application
Yaoyi Zhang	39900946	NPJ Sci Food	WB
Yuanyuan Zhang	39825194	Inflammation	WB
Yuchien Hsu	39807526	Adv Sci (Weinh)	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Non-treated and Calyculin A treated HeLa cells were subjected to SDS PAGE followed by western blot with 82349-1-RR (Phospho-IkB Alpha (Ser32/36) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-Ig) antibody as loading control.




Non-treated and Calyculin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 82349-1-RR (Phospho-IkB Alpha (Ser32/36) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-Ig) antibody as loading control.




Immunohistochemical analysis of paraffin-embedded skin cancer slide using 82349-1-RR (Phospho-IkB Alpha (Ser32/36) antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Non-treated and Calyculin A treated HSC-T6 cells were subjected to SDS PAGE followed by western blot with 82349-1-RR (Phospho-IkB Alpha (Ser32/36) antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with Alpha Tubulin (66031-1-Ig) antibody as loading control.



Immunohistochemical analysis of paraffin-embedded skin cancer slide using 82349-1-RR (Phospho-IkB Alpha (Ser32/36) antibody) at dilution of 1:2000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10⁶ HeLa cells untreated (dashed lines) or treated with Calyculin A which intracellularly stained with 0.06 ug Phospho-IkB Alpha (Ser32/36) Recombinant antibody (82349-1-RR, Clone:1G21) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.06 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with 90% MeOH.