

For Research Use Only

Phospho-p70(S6K) (Thr389) Recombinant antibody



Catalog Number: 82373-1-RR

Basic Information

Catalog Number: 82373-1-RR	GenBank Accession Number: NM_003161	Purification Method: Protein A purification
Size: 1000 µg/ml	GeneID (NCBI): 6198	CloneNo.: 1G15
Source: Rabbit	UNIPROT ID: P23443	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Full Name: ribosomal protein S6 kinase, 70kDa, polypeptide 1	
	Observed MW: 65-85 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : IGF-1 treated MCF-7 cells,
Species Specificity: Human	

Background Information

The Rps6kb1 gene encodes the 70 kDa ribosomal protein S6 kinase (p70S6K), which is a serine/threonine kinase regulated by phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin (mTOR) pathway. P70S6K plays a crucial role in controlling cell cycle, growth and survival. The PI3K/mTOR signalling pathway is one of the major mechanisms for controlling cell survival, proliferation and metabolism and is the central regulator of translation of some components of protein synthesis system. Due to alternative translation two isoform S6K1 proteins are known to exist in mammalian cells: p85 S6K1 and p70 S6K1, which is identical to p85 S6K but lacks its first 23 amino acids. In addition, mammalian cells express a second S6K1 isoform spanning 316 amino acids (p31 S6K1). mTOR is known to phosphorylate and thereby activate p70 S6K1 at T389 and p85 S6K1 at T412. (PMID: 25100792, PMID: 24970012, PMID: 21602892)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated and IGF-1 treated MCF-7 cells were subjected to SDS PAGE followed by western blot with 82373-1-RR (Phospho-p70(S6K) (Thr389) antibody) at dilution of 1:40000 incubated at room temperature for 1.5 hours. The membrane was stripped and re-blotted with beta actin antibody (66009-1-Ig) as loading control.