

For Research Use Only

EIF3D Recombinant antibody

Catalog Number: 82512-1-RR

Featured Product



Basic Information

Catalog Number:

82512-1-RR

Size:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0268

GenBank Accession Number:

BC000328

GeneID (NCBI):

8664

UNIPROT ID:

O15371

Full Name:

eukaryotic translation initiation factor 3, subunit D

Calculated MW:

66 kDa

Observed MW:

65 kDa

Purification Method:

Protein A purification

CloneNo.:

308

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:1000-1:4000

IF/ICC 1:500-1:2000

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Species Specificity:

human, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, rat liver tissue, HEK-293 cells, HepG2 cells, A549 cells, A431 cells, Jurkat cells

IP : A549 cells,

IHC : human breast cancer tissue,

IF/ICC : U2OS cells,

Background Information

The mammalian translation initiation factor 3 (eIF3), is a multiprotein complex of ~600 kDa that binds to the 40 S ribosome and promotes the binding of methionyl-tRNA_i and mRNA. The eIF357(p66) is the major RNA binding subunit in this complex. Human eIF3-p66 shares 64% sequence identity with a hypothetical *Caenorhabditis elegans* protein, presumably the p66 homolog. Deletion analyses of recombinant derivatives of eIF3-p66 show that the RNA-binding domain lies within an N-terminal 71-amino acid region rich in lysine and arginine.

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82512-1-RR (EIF3D antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.




rat liver tissue were subjected to SDS PAGE followed by western blot with 82512-1-RR (EIF3D antibody) at dilution of 1:30300 incubated at room temperature for 1.5 hours.




WB result of EIF3D antibody (82512-1-RR; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EIF3D transfected HEK-293 cells.




IP result of anti-EIF3D (IP:82512-1-RR, 4ug; Detection:82512-1-RR 1:5000) with A549 cells lysate 920 ug.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 82512-1-RR (EIF3D antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 82512-1-RR (EIF3D antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0).



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using EIF3D antibody (82512-1-RR, Clone: 308) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed U2OS cells using EIF3D antibody (82512-1-RR, Clone: 308) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002).



Bi-layer interferometry (BLI) kinetic assays of 82512-1-RR against Human EIF3D were performed. The affinity constant is 5.83 nM.