For Research Use Only

Stathmin 1 Recombinant antibody

Catalog Number:82559-1-RR

Featured Product




Catalog Number: GenBank Accession Number: **Purification Method: Basic Information** 82559-1-RR BC014353 Protein A purification GenelD (NCBI): Concentration: CloneNo.: 1000 ug/ml 3925 2H6 UNIPROT ID: Recommended Dilutions: Source: Rabbit P16949 WB 1:2000-1:10000 IHC 1:1000-1:4000 Full Name: Isotype: IF-P 1:250-1:1000 lgG stathmin 1/oncoprotein 18 IF/ICC 1:500-1:2000 Calculated MW: Immunogen Catalog Number: AG1633 18 kDa **Observed MW:** 18 kDa **Tested Applications:** Positive Controls: **Applications** WB, IHC, IF/ICC, IF-P, ELISA WB: SH-SY5Y cells, HepG2 cells, Jurkat cells, PC-12 **Species Specificity:** cells human, mouse, rat IHC : rat brain tissue, human tonsillitis tissue Note-IHC: suggested antigen retrieval with IF-P: mouse brain tissue, TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate IF/ICC : HeLa cells, NIH/3T3 cells buffer pH 6.0 **Background Information** Stathmin 1 (STMN1) normally regulates microtubule dynamics either by sequestering free tubulin heterodimers or by promoting microtubule catastrophe. STMN1 is highly expressed in fetal and adult brain, spinal cord, and cerebellum. Many different phosphorylated forms are observed depending on specific combinations among the sites which can be phosphorylated. Phosphorylation of stathmin is involved in response to NGF, neuron polarization and microtubule polymerization inhibition activity. Increased expression of STMN1 has been observed in a variety of human malignancies, such as colorectal primary tumors and metastatic tissues, but its association with melanoma is so far not well known. Storage: Storage Store at -20°C. Stable for one year after shipment. Storage Buffer PBS with 0.02% sodium azide and 50% glycerol, pH7.3 Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82559-1-RR (Stathmin 1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of Stathmin 1 antibody (82559-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-Stathmin 1 transfected HepG2 cells.




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 82559-1-RR (Stathmin 1 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded rat brain tissue slide using 82559-1-RR (Stathmin 1 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 82559-1-RR (Stathmin 1 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse brain tissue using Stathmin 1 antibody (82559-1-RR, Clone: 2H6) at dilution of 1:500 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using Stathmin 1 antibody (82559-1-RR, Clone: 2H6) at dilution of 1:1000 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Immunofluorescent analysis of (-20°C Methanol) fixed NIH/3T3 cells using Stathmin 1 antibody (82559-1-RR, Clone: 2H6) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Biolayer interferometry (BLL) kinetic assays of 82559-1-RR against Human Stathmin 1 were performed. The affinity constant is 7.57 nM.