For Research Use Only

CDX2 Recombinant antibody

Catalog Number:82659-1-RR





Basic Information	Catalog Number: 82659-1-RR	GenBank Accession Number: BC014461	Purification Method: Protein A purification	
	Size:	GenelD (NCBI):	CloneNo.:	
	500 µg/ml	1045	3]1	
	Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG17310	UNIPROT ID: Q99626 Full Name: caudal type homeobox 2	Recommended Dilutions: WB 1:1000-1:5000 IF 1:550-1:2200	
				Calculated MW: 313 aa, 34 kDa Observed MW: 33-40 kDa
	Applications	Tested Applications:	Positive C	ontrols:
IF/ICC, WB, ELISA		WB : Caco-2 cells, mouse colon tissue, COLO 320 cells IF : Caco-2 cells,		
Species Specificity:				
	Human, mouse			
Background Informatio	CDX2, also named as Homeobox p binding domain and belongs to th transcriptional regulation of multi CDX1 to CDX2 protein may be imp defining patterns of proliferation expressed to reduce tumorigenic	protein CDX-2, is a 313 amino acid pro e Caudal homeobox family. CDX2 loc iple genes expression in the intestina portant in the anterior to posterior patt	tein, which contains one homeobox DNA- alizes in the nucleus and is involved in the lepithelium. The relative expression of erning of the intestinal epithelium and in lus axis. Both Cdx1 and Cdx2 genes must be optosis, and to reduce cell migration,	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





mouse colon tissue were subjected to SDS PAGE followed by western blot with 82659-1-RR (CDX2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed Caco-2 cells using CDX2 antibody (82659-1-RR, Clone: 3J1) at dilution of 1:1100 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).

Various lysates were subjected to SDS PAGE followed by western blot with 82659-1-RR (CDX2 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.