For Research Use Only

P21 Recombinant antibody

Catalog Number:82669-2-RR 1 Publications



Basic Information

Applications

Catalog Number: 82669-2-RR Concentration:

1000 ug/ml Source: Rabbit Isotype:

Immunogen Catalog Number:

AG0368

Tested Applications: WB, IF/ICC, IP, ELISA

Cited Applications:

Species Specificity:

human Cited Species: human

GenBank Accession Number:

BC000275 GeneID (NCBI): 1026

UNIPROT ID: P38936 Full Name:

(p21, Cip1)

cyclin-dependent kinase inhibitor 1A protein lysate IF/ICC 1:1000-1:4000

Calculated MW: 18 kDa Observed MW: 21 kDa

Positive Controls:

WB: A549 cells, MCF-7 cells, HCT 116 cells, HUVEC cells, HeLa cells, HepG2 cells, THP-1 cells, U-87 MG

Purification Method:

Protein A purification

Recommended Dilutions:

IP 0.5-4.0 ug for 1.0-3.0 mg of total

WB 1:500-1:3000

CloneNo.:

3N14

cells

IP: MCF-7 cells, IF/ICC: MCF-7 cells,

Background Information

CDKN1A (p21, CIP1, WAF1) is a cyclin-dependent kinase inhibitor. CDKN1A binds to and inhibits the activity of $cyclin-CDK2 \ or \ -CDK4 \ complexes, and thus functions \ as \ a \ regulator \ of \ cell \ cycle \ progression \ at \ the \ G1 \ phase. The$ expression of CDKN1A is induced by wild-type but not mutant p53 protein, through which CDKN1A mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. CDKN1A can interact with proliferating cell nuclear antigen (PCNA), and plays a regulatory role in S phase DNA replication and DNA damage repair. CDKN1A was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of CDK2, and may be instrumental in the execution of apoptosis following caspase activation. Two alternatively spliced variants, which encode an identical protein, have been reported.

Notable Publications


Author	Pubmed ID	Journal	Application
Neng Li	38309505	J Biol Chem	WB

Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82669-2-RR (P21 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using P21 antibody (82669-2-RR, Clone: 3N14) at dilution of 1:2000 and Coralite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), CL594-Phalloidin (red).



IP result of anti-P21 (IP:82669-2-RR, 4ug; Detection:82669-2-RR 1:800) with MCF-7 cells lysate 760 ug.



Biolayer interferometry (BLL) kinetic assays of 82669-2-RR against Human P21 were performed. The affinity constant is 112 nM.