For Research Use Only

# PD-L1/CD274 Recombinant antibody

Catalog Number:82719-15-RR

proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

## Basic Information

Catalog Number:
82719-15-RR

Size:
1000 μg/ml

Source:
Rabbit

Isotype:
IgG

Immunogen Catalog Number:
AG12432

GenBank Accession Number:
BC074984

GeneID (NCBI):
29126

UNIPROT ID:
Q9NZQ7

Full Name:
CD274 molecule

Calculated MW:
290 aa, 33 kDa

Observed MW:
50 kDa

Purification Method:
Protein A purification

CloneNo.:
2H4

Recommended Dilutions:
WB 1:1000-1:4000

## Applications

Tested Applications:
WB, ELISA

Species Specificity:
Human

Positive Controls:

WB : MDA-MB-231 cells, U-87 MG cells, human placenta tissue

## Background Information

Programmed cell death ligand 1 (PD-L1, CD274, or B7-H1), is the first member of B7 family to be discovered. B7 family molecules are type I transmembrane proteins belonging to the immunoglobulin superfamily. In concert with their CD28 family receptors, the B7s are key regulators of the adaptive immune response. PD-L1 is suggested as a negative regulator of T and B cell, and plays important role in mediating tolerance of lymphocytes to self-antigens. It is also involved in the costimulatory signal, essential for T-cell proliferation and production of IL10 and IFNG, in an IL2-dependent and a PD-1-independent manner. PD-L1 is a 290 aa transmembrane protein with a calculated molecular weight of 33 kDa, it is predicted to be 27-30 kDa after signal peptide cleavage (PMID: 25609200; 17076679). The apparent molecular weight has also been reported as 45-70 kDa, major glycosylated form of 45-50 kDa and multiple post-translational modifications form of 65-70 kDa (PMID: 18760278; 16493058).

## Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82719-15-RR (PD-L1/CD274 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.