

For Research Use Only

# CD79A Recombinant antibody

Catalog Number: 82747-6-RR



## Basic Information

<b>Catalog Number:</b> 82747-6-RR	<b>GenBank Accession Number:</b> BC113733	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 1000 µg/ml	<b>GeneID (NCBI):</b> 973	<b>CloneNo.:</b> 230167A8
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P11912	<b>Recommended Dilutions:</b> WB 1:5000-1:50000
<b>Isotype:</b> IgG	<b>Full Name:</b> CD79a molecule, immunoglobulin-associated alpha	
<b>Immunogen Catalog Number:</b> AG17924	<b>Calculated MW:</b> 226 aa, 25 kDa	
	<b>Observed MW:</b> 38-45 kDa	

## Applications

<b>Tested Applications:</b> WB, ELISA	<b>Positive Controls:</b> WB : Daudi cells, Ramos cells, Raji cells
<b>Species Specificity:</b> Human	

## Background Information

CD79A, also named as B-cell antigen receptor complex-associated protein alpha chain or MB-1 membrane glycoprotein, is a 226 amino acid protein, which contains one ITAM domain and one Ig-like C2-type (immunoglobulin-like) domain. CD79A is expressed in B cells and localizes in the cell membrane. CD79A is required in cooperation with CD79B for initiation of the signal transduction cascade activated by binding of antigen to the B-cell antigen receptor complex (BCR) which leads to internalization of the complex, trafficking to late endosomes and antigen presentation. CD79A is also required for BCR surface expression and for efficient differentiation of pro- and pre-B-cells.

## Storage

**Storage:**  
Store at -20°C. Stable for one year after shipment.  
**Storage Buffer:**  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)

W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82747-6-RR (CD79A antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.