

For Research Use Only

Phospho-ULK1 (Ser758) Recombinant monoclonal antibody, PBS Only

Catalog Number: 82776-8-PBS



Basic Information

Catalog Number:

82776-8-PBS

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_003565

GeneID (NCBI):

8408

UNIPROT ID:

O75385

Full Name:

unc-51-like kinase 1 (C. elegans)

Observed MW:

140 kDa

Purification Method:

Protein A purification

CloneNo.:

240770A7

Applications

Tested Applications:

WB, Indirect ELISA

Species Specificity:

human

Background Information

Unc-51-like-kinase 1 (ULK1) is a target of both the mechanistic target of rapamycin (mTOR) and AMP activated protein kinase (AMPK), whose role is to facilitate the initiation of autophagy in response to starvation. ULK1 is phosphorylated on serine 638 and 758 sites by mTOR in nutrient-rich conditions, inhibiting ULK1 activation by disrupting its binding to AMPK. Upon glucose starvation, dissociation of mTOR from ULK1 and phosphorylation by AMPK leads to the activation of ULK1 activity. The S556 site of ULK1 is one of the major AMPK-dependent phosphorylation sites. (PMID: 30517873, PMID: 21258367). This antibody is equivalent to Ser757 in mice.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS only, pH7.3

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Non-treated and IGF-1 treated MCF-7 cells were subjected to SDS PAGE followed by western blot with 82776-8-RR (Phospho-ULK1 (Ser758) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was tripped and re-blotted with Alpha Tubulin antibody (66031-1-Ig) as loading control. This data was developed using the same antibody clone with 82776-8-PBS in a different storage buffer formulation.



Non-treated, phosphatase inhibitor treated and phosphatase treated PC-3 cells were subjected to SDS PAGE followed by western blot with 82776-8-RR (Phospho-ULK1 (Ser758) antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was tripped and re-blotted with Vinculin antibody (66305-1-Ig) as loading control. This data was developed using the same antibody clone with 82776-8-PBS in a different storage buffer formulation.