

# CD55 Recombinant antibody

Catalog Number: 82781-6-RR

## Basic Information

<b>Catalog Number:</b> 82781-6-RR	<b>GenBank Accession Number:</b> BC001288	<b>Purification Method:</b> Protein A purification
<b>Size:</b> 500 µg/ml	<b>GeneID (NCBI):</b> 1604	<b>CloneNo.:</b> 2D2
<b>Source:</b> Rabbit	<b>UNIPROT ID:</b> P08174	<b>Recommended Dilutions:</b> WB 1:2000-1:10000 IHC 1:200-1:800
<b>Isotype:</b> IgG	<b>Full Name:</b> CD55 molecule, decay accelerating factor for complement (Cromer blood group)	
<b>Immunogen Catalog Number:</b> AG25026	<b>Calculated MW:</b> 41 kDa	
	<b>Observed MW:</b> 75 kDa	

## Applications

**Tested Applications:**

IHC, WB, ELISA

**Species Specificity:**

Human

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

**Positive Controls:**

**WB:** HeLa cells, A549 cells, MKN-45 cells, human placenta tissue, K-562 cells

**IHC:** human lung cancer tissue, human placenta tissue

## Background Information

CD55, also known as DAF, is a glycosylphosphatidylinositol (GPI)-anchored surface glycoprotein that is widely distributed on blood, stroma, epithelial, and endothelial cells (PMID: 7517044; 29503741). It can also exist as a soluble form in plasma, urine, saliva, tears, and synovial fluids (PMID: 29503741). CD55 is a complement regulatory protein (PMID: 2469439; 7517044). It inhibits formation of the C3 convertases through binding to C3b and C4b. It also binds the alternate pathway convertase C3bBb, the classical pathway convertase and C4b2a to accelerate their decay (PMID: 17289551). CD55 also serves as a receptor for coxsackieviruses B1, B3, and B5 and several enteroviruses (PMID: 7538177; 7517044). The observed molecular weight of mature CD55 varies between 50 to 100 kDa depending on the cell type. Different sizes of CD55 might be caused by alternative splicing or different glycosylation patterns (PMID: 29503741).

## Storage

**Storage:**

Store at -20°C. Stable for one year after shipment.

**Storage Buffer:**

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82781-6-RR (CD55 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 82781-6-RR (CD55 antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 82781-6-RR (CD55 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).