For Research Use Only

SIRT1 Recombinant antibody, PBS Only




Catalog Number:82790-4-PBS

Basic Information	Catalog Number: 82790-4-PBS	GenBank Accession Number: BC012499	Purification Method: Protein A purification
	Size: 1mg/ml	GenelD (NCBI): 23411	CloneNo.: 2F11
	Source: Rabbit	UNIPROT ID: Q96EB6	
	Isotype: IgG Immunogen Catalog Number: AG3808	Full Name: sirtuin (silent mating type information regulation 2 homolog) 1 (S. cerevisiae)	
		Calculated MW: 747 aa, 82 kDa	
		Observed MW: 110-130 kDa	
Applications	Tested Applications: WB, IP, IF/ICC, Indirect ELISA Species Specificity:		
	Human		
Background Information	SIRT1, also named as SIR2L1, contains a deacetylase sirtuin-type domain and belongs to the sirtuin family. The post- translation modified SIRT1 is a 110-130 kDa protein, which contains one deacetylase sirtuin-type domain. The 75-80 kDa SirT1 fragment was detected to lack the carboxy-terminus (PMID:21305533). SirT1 exists a 57-61 kDa isoform. SIRT1 may be found in nucleolus, nuclear euchromatin, heterochromatin, and inner membrane. It can shuttles between the nucleus and cytoplasm. SIRT1 regulates processes such as apoptosis and muscle differentiation by deacetylating key proteins. SIRT1 in particular initiates several signaling events relevant to cardioprotection, including activation of endothelial nitric oxide synthase, INS receptor signaling, and autophagy. In addition, SIRT1 activation elicits resistance to oxidative stress via the regulation of transcription factors and co-activators such as FOXO, Hif-2a, and NF-kB. SIRT1 regulates the p53-dependent DNA damage response pathway by binding to and deacetylating p53, specifically at Lysine 382.		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer: PBS Only	cks. Upon receipt, store it immediately at	t -80°C


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82790-4-RR (SIRT1 antibody) at dilution of 1:4700 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SIRT1 antibody (82790-4-RR, Clone: 2F11) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 594-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR004). This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



IP result of anti-SIRT1 (IP:82790-4-RR, 4ug; Detection:82790-4-RR 1:1000) with HEK-293 cells lysate 1575 ug. This data was developed using the same antibody clone with 82790-4-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 82790-4-RR against Human SIRT1 were performed. The affinity constant is 4.90 nM.