For Research Use Only

IL-10 Recombinant antibody

Catalog Number:82793-16-RR 1 Publications



Purification Method:

Protein A purification

Recommended Dilutions:

WB 1:1000-1:4000

CloneNo.:

Basic Information

Catalog Number: 82793-16-RR

Size: 300 µg/ml Source: Rabbit

Isotype: IgG GenBank Accession Number:

NM_012854
GeneID (NCBI):
25325
UNIPROT ID:
P29456
Full Name:
interleukin 10

Calculated MW: 20kd Observed MW:

30 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications: WB

Species Specificity: rat, mouse Cited Species: mouse Positive Controls:

WB: Recombinant protein,

Background Information

The human IL10 gene is 4.7 kb in size and is located on the long arm of chromosome 1. It contains five exons. Biologically functional IL-10 exists in the form of a 36 kDa homodimer composed of two non-covalently bonded mono-mers each with 160 amino acids length. IL-10 is an anti-inflammatory cytokine, produced by T helper (Th) cells, macrophages, monocytes, and B cells, that plays a crucial role in preventing inflammatory and autoimmune pathologies. It downregulates the expression of Th1 cytokines, MHC class II antigens, and co-stimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. IL-10 can block NF- K B activity, and is involved in the regulation of the JAK-STAT signaling pathway. IL-10, along with its receptors, describes an important role in pathogenesis of various diseases, including infectious, inflammatory, autoimmune diseases. IL-10 mutations are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Notable Publications

Author	Pubmed ID	Journal	Application
Yeshen Zhang	39415256	J Nanobiotechnology	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



Recombinant protein were subjected to SDS PAGE followed by western blot with 82793-16-RR (IL-10 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.