

For Research Use Only

ICAM-1/CD54 Recombinant antibody

Catalog Number: 82827-8-RR



Basic Information

Catalog Number: 82827-8-RR	GenBank Accession Number: NM_010493	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 15894	CloneNo.: 240249H6
Source: Rabbit	UNIPROT ID: P13597	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF-P 1:250-1:1000
Isotype: IgG	Full Name: intercellular adhesion molecule 1	
	Calculated MW: 59 kDa	
	Observed MW: 85-110 kDa	

Applications

Tested Applications: WB, IHC, IF-P, ELISA	Positive Controls:
Species Specificity: mouse	WB : mouse kidney tissue, mouse lung tissue, mouse brain tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	IHC : mouse liver tissue,
	IF-P : mouse liver tissue,

Background Information

ICAM-1 (CD54) is a transmembrane glycoprotein of the immunoglobulin superfamily and is critical for the firm attachment and transmigration of leukocytes out of blood vessels and into tissues (PMID: 19307690). ICAM-1 is expressed by several cell types, typically on endothelial cells and cells of the immune system, and its expression can be up-regulated by various stimuli, including TNF- α , INF- γ , IL-1 and thrombin (PMID: 3086451; 9694714; 15979056). It is a ligand for LFA-1 and Mac-1, serves as a receptor for rhinovirus, and is one of several receptors used by Plasmodium falciparum (PMID: 2566624; 2538244; 2475784).


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82827-8-RR (ICAM1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 82827-8-RR (ICAM-1/CD54 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 82827-8-RR (ICAM-1/CD54 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse liver tissue using ICAM-1/CD54 antibody (82827-8-RR, Clone: 240249H6) at dilution of 1:500 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 82827-8-RR against Mouse ICAM-1 were performed. The affinity constant is 0.714 nM.