

For Research Use Only

# Acetyl-Histone H3 (Lys18) Recombinant antibody

Catalog Number: 82832-1-RR



## Basic Information

Catalog Number:

82832-1-RR

Concentration:

1000 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

BC066245

GeneID (NCBI):

8350

UNIPROT ID:

P68431

Full Name:

histone cluster 1, H3a

Observed MW:

15 kDa

Purification Method:

Protein A purification

CloneNo.:

2E3

Recommended Dilutions:

WB 1:5000-1:50000

## Applications

Tested Applications:

WB, Dot Blot, ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB : Trichostatin A treated NIH/3T3 cells, HSC-T6 cells, NIH/3T3 cells

## Background Information

Histones, including H1/H5 (linker histones), H2, H3, and H4 (core histones), are nucleic proteins which interact with DNA to form the nucleosomes and play important roles in gene regulation and DNA replication. Histone proteins are highly post-translationally modified while Histone H3 is the most extensively modified.

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Dot blot analysis was used to confirm the specificity of 82832-1-RR Acetyl-Histone H3 (Lys18) antibody. Acetylated peptides were spotted onto NC and probed with antibody at 1 µg/mL. The amount of peptide (µg/mL) spotted is indicated next to each row. Column 1: H3AK18Ac. Column 2: Unmodified H3AK18. Column 3: H3AK36Ac. Column 4: Unmodified H3AK36. Column 5: H3AK23Ac. Column 6: Unmodified H3AK23. Column 7: H3AK122Ac. Column 8: Unmodified H3AK122. Column 9:



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 82832-1-RR (Acetyl-Histone H3 (Lys18) antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Trichostatin A treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 82832-1-RR (Acetyl-Histone H3 (Lys18) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with Alpha Tubulin Monoclonal antibody (66031-1-Ig) as loading control.



HSC-T6 cells were subjected to SDS PAGE followed by western blot with 82832-1-RR (Acetyl-Histone H3 (Lys18) antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.