

For Research Use Only

Lipocalin-2/NGAL Recombinant antibody

Catalog Number: 82841-1-RR

Featured Product



Basic Information

Catalog Number:

82841-1-RR

Size:

500 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NP_005555.2

GeneID (NCBI):

3934

UNIPROT ID:

P80188

Full Name:

lipocalin 2

Calculated MW:

23kd

Observed MW:

22-25 kDa

Purification Method:

Protein A purification

CloneNo.:

2G1

Recommended Dilutions:

WB 1:2000-1:10000

IHC 1:50-1:500

Applications

Tested Applications:

WB, IHC, ELISA

Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : human saliva, HT-29 cells, SW480 cells

IHC : human intrahepatic cholangiocarcinoma tissue,

Background Information

Lipocalin-2 (LCN-2) is a adipocytokine also referred to as neutrophil gelatinase-associated lipocalin (NGAL). Lipocalin-2 is a circulatory protein responsible for the transportation of small and hydrophobic molecules to target organs. Lipocalin-2 is used as a biomarker for acute and chronic renal injury. It is present in a large variety of cells including neutrophil, hepatocytes, lung, bone marrow, adipose tissue, macrophages, thymus, non-neoplastic breast duct, prostate, and renal cells. Different functions have been associated with Lipocalin-2, including antibacterial, anti-inflammatory, and protection against cell and tissue stress (PMID:34463264).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82841-1-RR (Lipocalin-2/NGAL antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



WB result of NGAL antibody (82841-1-RR; 1:400; incubated at room temperature for 1.5 hours) with sh-Control and sh-NGAL transfected HT-29 cells.



Immunohistochemical analysis of paraffin-embedded human intrahepatic cholangiocarcinoma tissue slide using 82841-1-RR (NGAL antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 82841-1-RR against Human NGAL were performed. The affinity constant is 0.776 nM.