

For Research Use Only

Lipocalin-2/NGAL Recombinant antibody, PBS Only (Capture)

Catalog Number: 82841-9-PBS



Basic Information

Catalog Number:

82841-9-PBS

Size:

1mg/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NP_005555.2

GeneID (NCBI):

3934

UNIPROT ID:

P80188

Full Name:

lipocalin 2

Calculated MW:

23kd

Purification Method:

Protein A purification

CloneNo.:

2N10

Applications

Tested Applications:

Sandwich ELISA, Indirect ELISA, Sample test

Species Specificity:

human

Background Information

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Sandwich ELISA standard curve of MP00770-1, NGAL Recombinant Matched Antibody Pair - PBS only. 82841-9-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Eg0268. 82841-2-PBS was HRP conjugated as the detection antibody. Range: 19.5-1250 pg/mL




Human peripheral blood mononuclear cells (PBMC) were cultured unstimulated or stimulated with 10 μ g/mL PHA for 1 days. The mean NGAL concentration was determined to be 18,080.8 pg/mL in unstimulated PBMC supernatant, 32,268.8 pg/mL in PHA stimulated PBMC supernatant.



Saliva of eight individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 158,732.5 pg/mL with a range of 52,136.9 - 388,400.2 pg/mL.



Urine of eight individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 11,654.6 pg/mL with a range of 1,964.2 - 21,150.4 pg/mL



Plasma of fifteen individual healthy human donors was measured. The NGAL concentration of detected samples was determined to be 38,260.8 pg/mL with a range of 17,778.7 - 90,593.8 pg/mL