

For Research Use Only

RBM17 Recombinant antibody, PBS Only

Catalog Number: 82889-1-PBS

Featured Product



Basic Information

Catalog Number:

82889-1-PBS

Size:

1mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG0985

GenBank Accession Number:

BC007871

GeneID (NCBI):

84991

UNIPROT ID:

Q96125

Full Name:

RNA binding motif protein 17

Calculated MW:

45 kDa

Observed MW:

45-50 kDa

Purification Method:

Protein A purification

CloneNo.:

230125F6

Applications

Tested Applications:

WB, IF/ICC, IP, Indirect ELISA

Species Specificity:

human

Background Information

RNA-binding motif protein 17 (RBM17), also named splicing factor 45, was first identified as a member of the spliceosome. RBM17 expression is low in normal tissues but is overexpressed in several forms of cancer, including breast, ovarian and prostate.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using RBM17 antibody (82889-1-RR, Clone: 230125F6) at dilution of 1:1000 and Multi-Ab CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004).(CL488-Phalloidin, green). This data was developed using the same antibody clone with 82889-1-PBS in a different storage buffer formulation.




WB result of RBM17 antibody (82889-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-RBM17 transfected MCF-7 cells. This data was developed using the same antibody clone with 82889-1-PBS in a different storage buffer formulation.



IP result of anti-RBM17 (IP:82889-1-RR, 4ug; Detection:82889-1-RR 1:10000) with MCF-7 cells lysate 1470 ug. This data was developed using the same antibody clone with 82889-1-PBS in a different storage buffer formulation.



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug RBM17 Recombinant antibody (82889-1-RR, Clone:230125F6) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 82889-1-PBS in a



Biolayer interferometry (BLI) kinetic assays of 82889-1-RR against Human RBM17 were performed. The affinity constant is 1.54 nM.