For Research Use Only

MELK Recombinant antibody, PBS Only (Capture/Detector)

Catalog Number:82893-2-PBS Featured Product



Purification Method:

Protein A purification

CloneNo.:

230106A4

Basic Information

Catalog Number: 82893-2-PBS Concentration:

1mg/ml Source: Rabbit Isotype:

gG maternal embryonic leucine zipper

Immunogen Catalog Number: kinase

AG1897 Calculated MW:

651 aa, 75 kDa Observed MW: 50-70 kDa

BC014039

9833

Q14680 Full Name:

GeneID (NCBI):

UNIPROT ID:

GenBank Accession Number:

Applications

Tested Applications:

 $WB, IHC, Cytometric\,bead\,array, Sandwich\,ELISA,\\$

Indirect ELISA, Sample test

Species Specificity: human, mouse

Background Information

MELK is involved in various processes such as cell cycle regulation, self-renewal of stem cells, apoptosis and splicing regulation. It also plays a key role in cell proliferation and carcinogenesis, and it is required for proliferation of embryonic and postnatal multipotent neural progenitors.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: 100% PBS pH 7.3


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 82893-2-RR (MELK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 82893-2-PBS in a different storage buffer formulation.




WB result of MELK antibody (82893-2-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MELK transfected HeLa cells. This data was developed using the same antibody clone with 82893-2-PBS in a different storage buffer formulation.




Sandwich ELISA standard curve of MP00075-4, Human MELK Recombinant Matched Antibody Pair - PBS only. 82893-2-PBS was coated to a plate as the capture antibody and incubated with serial dilutions of standard Ag1897. 82893-6-PBS was HRP conjugated as the detection antibody. Range: 0.313-20 ng/mL




NCCIT and K-562 cell lysate were measured. The human MELK concentration of detected samples was determined to be 15.9 ng/mL (based on a 4.5 mg/mL extract load) in NCCIT cell lysate and 16.4 ng/mL (based on a 4.5 mg/mL extract load) in K-562 cell lysate.




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 82893-2-RR (MELK antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82893-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse kidney tissue slide using 82893-2-RR (MELK antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 82893-2-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP00075-1, MELK Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 82893-2-PBS. Detection antibody: 82893-3-PBS. Standard: Ag1897. Range: 0.313-20 ng/mL



Cytometric bead array standard curve of MP00075-2, MELK Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 82893-4-PBS. Detection antibody: 82893-2-PBS. Standard: Ag1897. Range: 0.313-20 ng/mL



Biolayer interferometry (BLI) kinetic assays of 82893-2-RR against Human MELK were performed. The affinity constant is below 1 pM.