## For Research Use Only

## PARK7,DJ-1 Recombinant antibody

Catalog Number:82913-1-RR



**Basic Information** 

Catalog Number:

GenBank Accession Number: BC008188

Purification Method:

82913-1-RR

\_\_\_\_\_

Protein A purification

Recommended Dilutions:

Size:

GeneID (NCBI):

CloneNo.:

1000 μg/ml

11315 UNIPROT ID: 230124E12

Source: Rabbit

Q99497

IP 0.5-4.0 ug for 1.0-3.0 mg of total

Isotype:

Full Name:

protein lysate

IgG

Parkinson disease (autosomal recessive, early onset) 7

IF 1:200-1:800

AG2287

Calculated MW: 189 aa, 20 kDa

**Applications** 

**Background Information** 

Tested Applications:

Immunogen Catalog Number:

Positive Controls:

IP, IF/ICC, FC, ELISA Species Specificity:

IP: HeLa cells,
IF: HepG2 cells,

Humar

PARK7, also named as DJ1, belongs to the peptidase C56 family. It protects cells against oxidative stress and cell death. PARK7 plays a role in regulating expression or stability of the mitochondrial uncoupling proteins SLC25A14 and SLC25A27 in dopaminergic neurons of the substantia nigra pars compacta and attenuates the oxidative stress induced by calcium entry into the neurons via L-type channels during pacemaking. It eliminates hydrogen peroxide and protects cells against hydrogen peroxide-induced cell death. PARK7 has cell-growth promoting activity and transforming activity. It may function as a redox-sensitive chaperone. It's precursor undergoes a cleavage of a C-terminal peptide and subsequent activation of protease activity in response to oxidative stress. The amino acid

replace at 166 (L  $\rightarrow$  P) reduces PARK7 protein stability and leads to increased degradation. The predicted MW of this protein is 20 kDa, An additional 25 kDa band can be observed due to modification (PMID: 31767755).

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




IP result of anti-PARK7,DJ-1 (IP:82913-1-RR, 4ug; Detection:82913-1-RR 1:2000) with HeLa cells lysate 1920 ug.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using PARK7,DJ-1 antibody (82913-1-RR, Clone: 230124E12) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug Anti-Human PARK7,DJ-1 (82913-1-RR, Clone:230124E12) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:166 (red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10^6 HepG2 cells were intracellularly stained with 0.2 ug Anti-Human PARk(7,D]-1 (82913-1-RR, Clone:230124E12) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:166 (red), or 0.2 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP, Clone:) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).