For Research Use Only

PICK1 Recombinant antibody

Catalog Number:82919-1-RR



Basic Information

Catalog Number:

GenBank Accession Number: BC017561

Purification Method: Protein A purification

82919-1-RR Size:

GeneID (NCBI):

CloneNo.:

800 μ g/ml

9463

230185B8

Source: Rabbit

UNIPROT ID: Q9NRD5

Recommended Dilutions: WB 1:1000-1:8000

Isotype:

AG1443

Full Name:

protein interacting with PRKCA 1

Immunogen Catalog Number:

Calculated MW:

47 kDa

Observed MW:

50 kDa

Applications

Tested Applications:

Positive Controls:

WB, FC, ELISA

Species Specificity: Human, mouse, rat, pig WB: LNCaP cells, HeLa cells, MCF-7 cells, mouse brain

tissue, rat brain tissue, pig brain tissue

Background Information

Protein interacting with C kinase 1 (PICK1) was first cloned as a PKC-binding partner through yeast two hybrid system. PICK1 acts as a critical regulator of membrane receptors' subcellular trafficking to modulate neural processes such as learning and memory, and is widely expressed in brain, testis, heart, lung, liver, kidney and muscle. It probably binds to and organize the subcellular localization of a variety of membrane proteins containing some PDZ recognition sequence, for instance, PICK1 is a critical mediator of α -amino-3-hydroxy-5-methyl-4isoxazolepropionic acid receptor (AMPAR) trafficking in neural synapses. PICK1 expression on D-serine release and glutamate transport in astrocytes suggests a potential implication of PICK1 in the progression of amyotrophic lateral sclerosis (ALS). PICK1 may also participate in breast cancer development through inhibition of TGF- β signaling.

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82919-1-RR (PICK1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 82919-1-RR (PICK1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



1x10^6 HEK-293T cells were intracellularly stained with 0.25 ug Anti-Human PICK1 (82919-1-RR, Clone:230185B8) and APC-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) at dilution 1:166 (red), or 0.25 ug Rabbit 1gG control Rabbit PolyAb (30000-0-AP, Clone:). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).