

For Research Use Only

RBP4 Recombinant antibody

Catalog Number: 82934-1-RR



Basic Information

Catalog Number: 82934-1-RR	GenBank Accession Number: BC020633	Purification Method: Protein A purification
Size: 400 µg/ml	GeneID (NCBI): 5950	CloneNo.: 6013
Source: Rabbit	UNIPROT ID: P02753	Recommended Dilutions: WB 1:1000-1:4000
Isotype: IgG	Full Name: retinol binding protein 4, plasma	
Immunogen Catalog Number: AG2448	Calculated MW: 201 aa, 23 kDa	
	Observed MW: 23 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls: WB : HepG2 cells,
Species Specificity: Human	

Background Information

RBP4 (retinol-binding protein 4) is a carrier protein that transports vitamin A (retinol) from the liver to the peripheral tissues. Synthesized primarily by hepatocytes and adipocytes as a 21 kDa non-glycosylated protein, RBP4 is secreted into the circulation as a retinol-RBP4 complex. In plasma the RBP4-retinol complex is bound to transthyretin (TRR), which prevents prevent kidney filtration. Two truncated forms of RBP4, RBP4-L (truncated at Leu-183) and RBP4-LL (truncated at Leu-182 and Leu-183), exist by proteolytic process. RBP4-L and RBP4-LL, which do not bind TTR, are normally excreted into the urine but accumulate in the serum during renal failure. Urinary RBP4 has been reported as marker for glomerular disease. RBP4 also was identified as an adipokine that elevated in some INS-resistant states. Measurement of serum RBP4 could be used to assess the risk of INS resistance, type 2 diabetes, obesity, and cardiovascular disease. (18752671, 16034410)

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com

W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 82934-1-RR (RBP4 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.