For Research Use Only

PAK7 Recombinant antibody

Catalog Number:82935-1-RR Featured Product





Basic Information	Catalog Number: 82935-1-RR	GenBank Accession Number: BC024179	Purification Method: Protein A purification	
	Concentration: 400 ug/ml	GenelD (NCBI): 57144	CloneNo.: 230201A10	
	Source: Rabbit	UNIPROT ID: Q9P286	Recommended Dilutions: WB 1:500-1:2000	
	Isotype: IgG Immunogen Catalog Number:	Full Name: p21 protein (Cdc42/Rac)-activated kinase 7 Calculated MW: 719 aa, 81 kDa		
	AG29190			
		Observed MW: 70 kDa		
Applications	Tested Applications: WB, FC (Intra), ELISA Species Specificity: human	Positive Controls:		
		WB : HeLa	cells, SH-SY5Y cells	
Background Information	found in the brain, which promote expression level of PAK7 significa hepatocellular carcinoma, pancre tumorigenesis, invasion, and met	PAK7, also known as PAK5, is a newly found but scarcely researched type II PAKs (PAK4, 5, 6). PAK7 was originally found in the brain, which promotes filopodia formation in nerve cells. Recently, studies have found that the expression level of PAK7 significantly increased in colorectal cancer, ovarian cancer, gastric cancer, neuroglioma, hepatocellular carcinoma, pancreatic cancer, osteosarcoma, and breast cancer. PAK7 plays an important role in tumorigenesis, invasion, and metastasis, mainly related to its mechanism of regulating cytoskeleton, inhibiting apoptosis, and promoting proliferation. The calculated molecular weight of PAK7 is 80 kDa (PMID: 29805709, PMID: 34165002).		
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20°	l 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact: E: Proteintech-CN@ptglab.com T: 4006900926 W: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





WB result of PAK7 antibody (82935-1-RR; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PAK7 transfected HeLa cells. Various lysates were subjected to SDS PAGE followed by western blot with 82935-1-RR (PAK7 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



(blue)1x10^6 K-562 cells were intracellularly stained with 0.25 ug Anti-Human PAK7 (82935-1-RR, Clone:230201A10) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) at dilution 1:166(red), or 0.25 ug Rabbit IgG control Rabbit PolyAb (30000-0-AP). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).