For Research Use Only

MMP13 Recombinant antibody, PBS Only Catalog Number:83188-4-PBS





Basic Information	Catalog Number: 83188-4-PBS	GenBank Accession Number: BC074808	Purification Method: Protein A purfication	
	Concentration: 1 mg/ml	GenelD (NCBI): 4322	CloneNo.: 240060A1	
	Source: Rabbit	UNIPROT ID: P45452		
	lsotype: IgG	Full Name: matrix metallopeptidase 13		
	Immunogen Catalog Number: AG34892	Calculated MW: 471 aa, 54 kDa		
		Observed MW: 70 kDa, 54 kDa		
Applications	Tested Applications: WB, FC (Intra), ELISA			
	Species Specificity: human			
Background Informat	analogue of human interstitial MM growth plate. The bands seen with reported for this enzyme, along w detected 3 bands that the glycosy	MMP-13(Matrix metalloproteinase-13), cleaves type I collagen and was previously thought to be the rodent analogue of human interstitial MMP-1. MMP-13 is found in hypertrophic and calcifying cartilage of mammalian growth plate. The bands seen with rat MMP-13 are consistent with a 59-kDa latent form and a 43-kDa active form reported for this enzyme, along with a smaller band at approximately 30 kDa(PMID:10771523). It also can be detected 3 bands that the glycosylated proMMP-13 (70 kDa), unglycosylated form (55 kDa), and active MMP-13 enzyme (40kDa) by western blot(PMID:18332263).		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer:	icks. Upon receipt, store it immediate	ly at −80°C	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data






Various lysates were subjected to SDS PAGE followed by western blot with 83188-4-RR (MMP13 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83188-4-PBS in a different storage buffer formulation. 1x10^6 MCF-7 cells were intracellularly stained with 0.25 ug MMP13 Recombinant antibody (83188-4-RR, Clone:240060A1) and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2)(red), or 0.25 ug Rabbit IgG Isotype Control Recombinant Antibody (98136-1-RR, Clone: 240953C9) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 83188-4-PBS in a

۰

▼



Biolayer interferometry (BLI) kinetic assays of 83188-4-RR against Human MMP13 were performed. The affinity constant is below 1 pM.