

For Research Use Only

CD163 Recombinant antibody

Catalog Number: 83285-4-RR



Basic Information

Catalog Number:

83285-4-RR

Size:

600 ug/ml

Source:

Rabbit

Isotype:

IgG

GenBank Accession Number:

NM_053094.2

GeneID (NCBI):

93671

UNIPROT ID:

Q2VLH6-1

Full Name:

CD163 antigen

Calculated MW:

121 kDa

Observed MW:

160-170 kDa

Purification Method:

Protein A purification

CloneNo.:

240233A3

Recommended Dilutions:

WB 1:2000-1:12000

IHC 1:2000-1:8000

IF-P 1:50-1:500

Applications

Tested Applications:

WB, IHC, IF-P, ELISA

Species Specificity:

mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : mouse liver tissue, mouse lung tissue, rat spleen tissue, rat liver tissue

IHC : mouse liver tissue, mouse spleen tissue

IF-P : mouse small intestine tissue, rat liver tissue

Background Information

CD163, also known as M130, is a membrane glycoprotein that belongs to the scavenger receptor superfamily (PMID: 8370408). It is an acute phase-regulated and signal-inducing macrophage protein expressed exclusively in monocytes and tissue macrophages (PMID: 11196644). CD163 mediates endocytosis of haptoglobin-haemoglobin complexes (PMID: 11196644). The uptake of haptoglobin by macrophages contributes to the recycling of iron and also to the inflammatory response (PMID: 22900885). Soluble CD163 (sCD163), as a result of ectodomain shedding during inflammatory activation of macrophages, circulates in blood and has been suggested as a plasma/serum marker for macrophage activity (PMID: 12570164).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83285-4-RR (CD163 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 83285-4-RR (CD163 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 83285-4-RR (CD163 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse small intestine tissue using CD163 antibody (83285-4-RR, Clone: 240233A3) at dilution of 1:200 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded rat liver tissue using CD163 antibody (83285-4-RR, Clone: 240233A3) at dilution of 1:200 and Rhodamine (TRITC)-conjugated Goat Anti-Rabbit IgG(H+L) (SA00007-2), CD20 antibody (31909-1-AP, Magenta), EPCAM/CD326 antibody (32189-1-AP, green). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 83285-4-RR against Mouse CD163 were performed. The affinity constant is 2.25 nM.