

SMARCA4 Recombinant antibody

Catalog Number: 83310-2-RR

Basic Information

Catalog Number: 83310-2-RR	GenBank Accession Number: BC150298	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 6597	CloneNo.: 240172F4
Source: Rabbit	UNIPROT ID: P51532	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:2000-1:8000 IF/ICC 1:125-1:500
Isotype: IgG	Full Name: SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4	
Immunogen Catalog Number: AG16256	Calculated MW: 1647 aa, 185 kDa	
	Observed MW: 185 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls: WB : HeLa cells, HepG2 cells, HSC-T6 cells, NIH/3T3 cells, HEK-293T cells, C6 cells IHC : human placenta tissue, mouse cerebellum tissue, mouse colon tissue, mouse lung tissue, rat cerebellum tissue, rat colon tissue, rat lung tissue IF/ICC : HepG2 cells,
Species Specificity: human, mouse, rat	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

SMARCA4, also named as BAF190A, BRG1, SNF2B and SNF2L4, belongs to the SNF2/RAD54 helicase family. SMARCA4 is a transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. It is a component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator complex. It is also involved in vitamin D-coupled transcription regulation via its association with the WINAC complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligand-bound VDR-mediated transrepression of the CYP27B1 gene.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83310-2-RR (SMARCA4 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with .




Immunohistochemical analysis of paraffin-embedded mouse colon tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with .




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).






Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat colon tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SMARCA4 antibody (83310-2-RR, Clone: 240172F4) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).

Immunohistochemical analysis of paraffin-embedded rat cerebellum tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffin-embedded rat lung tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




1x10⁶ Jurkat cells were intracellularly stained with 0.25 ug SMARCA4 Recombinant antibody (83310-2-RR, Clone:240172F4) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).

Immunohistochemical analysis of paraffin-embedded rat colon tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded rat lung tissue slide using 83310-2-RR (SMARCA4 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 83310-2-RR against Human SMARCA4 were performed. The affinity constant is 3.66 nM.