

# Osteopontin Recombinant monoclonal antibody

Catalog Number: 83341-1-RR

3 Publications

## Basic Information

## Catalog Number:

83341-1-RR

## Source:

Rabbit

## Isotype:

IgG

## Immunogen Catalog Number:

EG0754

## GenBank Accession Number:

BC007016

## GeneID (NCBI):

6696

## UNIPROT ID:

P10451-5

## Full Name:

secreted phosphoprotein 1

## Calculated MW:

314 aa, 35 kDa

## Observed MW:

60-66 kDa

## Purification Method:

Protein A purification

## CloneNo.:

240206A2

## Recommended Dilutions:

WB: 1:1000-1:6000

IHC: 1:200-1:800

IF/ICC: 1:125-1:500

FC (Intra): 0.25 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

## Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

## Cited Applications:

WB, IHC, IF

## Species Specificity:

human, mouse

## Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

## Positive Controls:

WB: MOLT-4 cells, HepG2 cells, HEK-293 cells

IHC: human cervical squamous cancer tissue,

IF/ICC: RAW 264.7 cells,

FC (Intra): U-937 cells, A549 cells

## Background Information

Osteopontin (OPN), also known as SPP1, is a secreted glycoprophosphoprotein that belongs to the small integrin-binding ligand N-linked glycoprotein (SIBLING) family. Originally isolated from bone, OPN has been found in kidneys, vascular tissues, biological fluids, and various tumor tissues (PMID: 15138464; 16406521). OPN can interact with CD44 and integrins and regulate diverse biological processes. It has a multifaceted role in bone development and remodeling, and is also involved in the inflammatory and immune response, oncogenesis and cancer progression. The very acidic nature of OPN, as well as the presence of variable posttranslational modifications, has led to anomalous migration in SDS-polyacrylamide gels and therefore to reports of different molecular weights for OPN (PMID: 8293561). Depending on the cell and tissue source and/or the SDS-PAGE system, OPN migrates with a molecular weight of 44-80 kDa, as well as at some smaller bands corresponding to peptide fragments (PMID: 8195113; 17890765).

## Notable Publications

Author	Pubmed ID	Journal	Application
Yueze Liu	40499532	Mol Carcinog	IHC
Yanna Wang	39954559	Tissue Cell	WB,IF
Chunmei Wen	39888412	J Gastroenterol	IF

## Storage

## Storage:

Store at -20°C. Stable for one year after shipment.

## Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83341-1-RR (OPN, SPP1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human cervical squamous cancer tissue slide using 83341-1-RR (Osteopontin antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed RAW 264.7 cells using OPN, SPP1 antibody (83341-1-RR, Clone: 240206A2) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



1x10<sup>6</sup> U-937 cells were intracellularly stained with 0.25 ug OPN, Spp1 Recombinant Antibody (83341-1-RR, Clone:240206A2) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.



1x10<sup>6</sup> A549 cells were intracellularly stained with 0.25 ug OPN, Spp1 Recombinant Antibody (83341-1-RR, Clone:240206A2) and PE-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.



Biolayer interferometry (BLI) kinetic assays of 83341-1-RR against Human OPN/SPP1 were performed. The affinity constant is 36.8 nM.