

For Research Use Only

# NEDD9 Recombinant antibody

Catalog Number: 83362-2-RR

Featured Product



## Basic Information

### Catalog Number:

83362-2-RR

### Concentration:

800 ug/ml

### Source:

Rabbit

### Isotype:

IgG

### Immunogen Catalog Number:

AG34170

### GenBank Accession Number:

BC040207

### GeneID (NCBI):

4739

### UNIPROT ID:

Q14511

### Full Name:

neural precursor cell expressed,  
developmentally down-regulated 9

### Calculated MW:

834 aa, 93 kDa

### Observed MW:

105 kDa, 115 kDa

### Purification Method:

Protein A purification

### CloneNo.:

240310F8

### Recommended Dilutions:

WB 1:2000-1:16000

IF/ICC 1:50-1:500

## Applications

### Tested Applications:

WB, IF/ICC, ELISA

### Species Specificity:

human

### Positive Controls:

WB : A549 cells, HepG2 cells, MCF-7 cells, U-251 cells,  
Raji cells, Jurkat cells

IF/ICC : MCF-7 cells, A549 cells

## Background Information

NEDD9, also known as CASL, HEF1, contains an N-terminal Src homology (SH) 3 domain and an adjacent domain composed of multiple SH2-binding motifs (PMID: 8668148). Cell cycle-regulated processing produces four isoforms: p115, p105, p65, and p55. Isoform p115 arises from p105 phosphorylation and appears later in the cell cycle. Isoform p55 arises from p105 as a result of cleavage at a caspase cleavage-related site and it appears specifically at mitosis. p105 and p115 are predominantly cytoplasmic and associated with focal adhesions, whereas p55 associates with the mitotic spindle (PMID: 9584194).

## Storage

### Storage:

Store at -20°C. Stable for one year after shipment.

### Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83362-2-RR (NEDD9 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 83362-2-RR (NEDD9 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of NEDD9 antibody (83362-2-RR; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-NEDD9 transfected HepG2 cells.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using NEDD9 antibody (83362-2-RR, Clone: 240310F8) at dilution of 1:200 and CoraLite® Plus 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Immunofluorescent analysis of (4% PFA) fixed A549 cells using NEDD9 antibody (83362-2-RR, Clone: 240310F8) at dilution of 1:200 and Multi-rAb CoraLite® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red).



Bilayer interferometry (BLI) kinetic assays of 83362-2 against Human NEDD9 were performed. The affinity constant is 0.173 nM.