

GenelD (NCBI):

UNIPROT ID: Q6NVY1

26275

CloneNo.:

240590A3

Concentration:

1 mg/ml


Source:

Rabbit


	Isotype: IgG Immunogen Catalog Number: AG6149	Full Name: 3-hydroxyisobutyryl-Coenzyme A hydrolase Calculated MW: 43 kDa Observed MW: 38 kDa
Applications	Tested Applications: WB, IF/ICC, FC (Intra), Cytometric bead array, Indirect ELISA Species Specificity:	
Background Information	human, rat HIBCH belongs to the enoyl-CoA hydratase/isomerase family. HIBCH has two isoforms with MW 43 kDa and 38 kDa. It is highly expressed in the liver and kidney, also detected in the heart, muscle and brain (at protein level), but not detected in the lung. Hydrolyzes 3-hydroxyisobutyryl-CoA (HIBYL-CoA), a saline catabolite. Has high activity toward isobutyryl-CoA. Could be an isobutyryl-CoA dehydrogenase that functions in valine catabolism. Also hydrolyzes 3- hydroxypropanoyl-CoA (PMID: 8824301).	
Storage	Storage: Store at -80°C.	s. Upon receipt, store it immediately at -80°C

For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83569-1-RR (HIBCH antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83569-1-PBS in a different storage buffer formulation.




WB result of HIBCH antibody (83569-1-RR; 1:20000; incubated at room temperature for 1.5 hours) with sh-Control and sh-HIBCH transfected HepG2 cells. This data was developed using the same antibody clone with 83569-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HIBCH antibody (83569-1-RR, Clone: 240590A3) at dilution of 1:200 and MultirAb CoraLite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002). This data was developed using the same antibody clone with 83569-1-PBS in a different storage buffer formulation.




1x10<sup>^6</sup> HepG2 cells were intracellularly stained with 0.25 ug HIBCH Recombinant antibody (83569-1-RR, Clone:240590A3) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 83569-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using HIBCH antibody (83569-1-RR, Clone: 240590A3) at dilution of 1:400 and MultirAb Coralite ® Plus 488-Goat Anti-Rabbit Recombinant Secondary Antibody (H+L) (RGAR002), CL594-phalloidin (red). This data was developed using the same antibody clone with 83569-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP00560-3, HIBCH Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 83569-3-PBS. Detection antibody: 83569-1-PBS. Standard: Ag6149. Range: 0.625-80 ng/mL



Biolayer interferometry (BLl) kinetic assays of 83569-1-RR against Human HIBCH were performed. The affinity constant is 1.37 nM.