For Research Use Only

ZBTB7A Recombinant antibody, PBS Only

Catalog Number:83616-5-PBS

Featured Product



Purification Method:

Protein A purfication

CloneNo.:

240555A6

Basic Information

Catalog Number: 83616-5-PBS

Size:
1 mg/ml
Source:
Rabbit
Isotype:

i i

> Calculated MW: 61 kDa Observed MW:

GenBank Accession Number:

zinc finger and BTB domain

BC113511

51341

O95365 Full Name:

GeneID (NCBI):

UNIPROT ID:

containing 7A

61 kDa, 75-80 kDa

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:

human

Background Information

ZBTB7A, also named as LRF, FBI 1, FBI -1, pokemon, ZBTB7, ZNF857A and DKFZp547O146, a member of the POK (POZ and Krüppel) family of transcription factors, plays a role in differentiation, oncogenesis, and adipogenesis (PMID: 35798238). ZBTB7A is involved in the instruction of early lymphoid progenitors to develop into B lineage by repressing T-cell instructive Notch signals. ZBTB7A is widely expressed in normal thymus and it is aberrantly overexpressed in human cancers (PMID: 28942243). ZBTB7A has 10 potential sumoylation sites located at lysine 61, 354, 371, 379, 383, 396, 486, 487, 536 and 539 (PMID: 20471975, 19471103). ZBTB7A has a predicted molecular weight of 61 kDa and the 75-80-kDa band could be the sumoylated form.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer PBS Only


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83616-5-RR (ZBTB7A antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83616-5-PBS in a different storage buffer formulation.




WB result of ZBTB7A antibody (83616-5-RR; 1:8000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ZBTB7A transfected HEK-293 cells. This data was developed using the same antibody clone with 83616-5-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ZBTB7A antibody (83616-5-RR, Clone: 240555A6) at dilution of 1:200 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 83616-5-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using ZBTB7A antibody (83616-5-RR, Clone: 240555A6) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gC(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 83616-5-PBS in a different storage buffer formulation.



1x10^6 HeLa cells were intracellularly stained with 0.25 ug ZBTB7A Recombinant antibody (83616-5-RR, Clone:240555A6) and APC-Conjugated AffiniPure Goat Anti-Rabbit I gG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 83616-5-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 83616-5-RR (ZBTB7A antibody) at dilution of 1:400 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer(pH9). This data was developed using the same antibody clone with 83616-5-PBS in a different storage buffer formulation.