For Research Use Only

OPTN/Optineurin Recombinant antibody

Catalog Number:83634-5-RR



Basic Information

Catalog Number: GenBank Accession Number: 83634-5-RR BC013876

GeneID (NCBI): Size: 1000 ug/ml 10133 **UNIPROT ID:** Source: Rabbit Q96CV9 Full Name: Isotype: optineurin Calculated MW: Immunogen Catalog Number:

AG1272 Observed MW:

66-70 kDa

66 kDa

Positive Controls:

WB: A431 cells, HeLa cells, human testis tissue

Purification Method:

Protein A purfication

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:200-1:800

CloneNo.:

240636A7

IHC: human gliomas tissue,

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA Species Specificity:

human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

OPTN, also named as FIP2, GLC1E, HIP7, HYPL, and NRP, plays a neuroprotective role in the eye and optic nerve. It is probably part of the TNF-alpha signaling pathway that can shift the equilibrium toward induction of cell death. OPTN may act by regulating membrane trafficking and cellular morphogenesis via a complex that contains Rab8 $and \ hungting tin \ (HD). \ OPTN \ may \ constitute \ a \ cellular \ target \ for \ adenovirus \ E3\ 14.7, \ an \ inhibitor \ of \ TNF-alpha$ functions, thereby affecting cell death. Defects in OPTN cause susceptibility to normal pressure glaucoma (NPG). OPTN mutated in adult-onset primary open angle glaucoma. It supports the protective role of OPTN in the trabecular meshwork.


Storage

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83634-5-RR (OPTN antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 83634-5-RR (OPTN antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human gliomas tissue slide using 83634-5-RR (OPTN antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 HeLa cells were intracellularly stained with 0.25 ug OPTN Recombinant antibody (83634-5-RR, Clone:240636A7) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(HL)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83634-5-RR against Human OPTN were performed. The affinity constant is 12.2 nM.