

For Research Use Only

DLAT Recombinant antibody

Catalog Number:83654-3-RR



Basic Information

Catalog Number: 83654-3-RR	GenBank Accession Number: BC039084	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 1737	CloneNo.: 240713C12
Source: Rabbit	UNIPROT ID: P10515	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:250-1:1000 IF/ICC 1:125-1:500
Isotype: IgG	Full Name: dihydrolipoamide S-acetyltransferase	
Immunogen Catalog Number: AG22831	Calculated MW: 647 aa, 70 kDa	
	Observed MW: 69-70 kDa	

Applications

Tested Applications:
WB, IHC, IF/ICC, FC (Intra), ELISA

Species Specificity:
human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HepG2 cells, Jurkat cells, MCF-7 cells, LNCaP cells, mouse brain tissue, rat brain tissue

IHC : human stomach cancer tissue,

IF/ICC : HepG2 cells,

Background Information

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83654-3-RR (DLAT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83654-3-RR (DLAT antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer(pH9).




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DLAT antibody (83654-3-RR, Clone: 240713C12) at dilution of 1:250 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10⁶ HepG2 cells were intracellularly stained with 0.25 ug DLAT Recombinant Antibody (83654-3-RR, Clone:240713C12) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



1x10⁶ HeLa cells were intracellularly stained with 0.25 ug DLAT Recombinant antibody (83654-3-RR, Clone:240713C12) and APC-Conjugated Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Rabbit IgG Isotype Control RecAb (98136-1-RR, Clone: 240953C9) (blue), or 0.25 ug DLAT Recombinant antibody (83654-3-RR, Clone: 240713C12) (green). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Bilayer interferometry (BLI) kinetic assays of 83654-3-RR against Human DLAT were performed. The affinity constant is 0.17 nM.