

For Research Use Only

FTCD Recombinant antibody

Catalog Number: 83664-2-RR



Basic Information

Catalog Number:

83664-2-RR

Size:

600 ug/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG16646

GenBank Accession Number:

BC136395

GeneID (NCBI):

10841

UNIPROT ID:

O95954

Full Name:

formiminotransferase

cyclodeaminase

Calculated MW:

541 aa, 59 kDa

Observed MW:

58 kDa

Purification Method:

Protein A purification

CloneNo.:

240654G9

Recommended Dilutions:

WB 1:2000-1:12000

Applications

Tested Applications:

WB, ELISA

Species Specificity:

human, mouse, rat

Positive Controls:

WB : mouse liver tissue, rat liver tissue, human kidney tissue

Background Information

FTCD (formimidoyltransferase cyclodeaminase) is critical for the catabolism of histidine, a . This process is one-carbon units that can enter the one-carbon/folate cycle, which provides methyl groups for arsenic metabolism (PMID: 30893314). In addition, FTCD could serve as a predictive and prognostic marker for patients with HCC (PMID: 37675273).

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83664-2-RR (FTCD antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours.



1×10^6 HepG2 cells were intracellularly stained with 0.25 μ g FTCD Recombinant antibody (83664-2-RR, Clone:240654G9) and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2) (red), or 0.25 μ g Rabbit IgG control Rabbit PolyAb (30000-O-AP) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Biolayer interferometry (BLI) kinetic assays of 83664-2-RR against Human FTCD were performed. The affinity constant is 6.37 nM.