

For Research Use Only

# UQCRC2 Recombinant antibody

Catalog Number: 83667-2-RR



## Basic Information

Catalog Number: 83667-2-RR	GenBank Accession Number: BC000484	Purification Method: Protein A purification
Size: 1000 ug/ml	GeneID (NCBI): 7385	CloneNo.: 240454D9
Source: Rabbit	UNIPROT ID: P22695	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:50-1:500
Isotype: IgG	Full Name: ubiquinol-cytochrome c reductase core protein II	
Immunogen Catalog Number: AG6432	Calculated MW: 48 kDa	
	Observed MW: 48 kDa	

## Applications

Tested Applications: WB, IHC, FC (Intra), ELISA	Positive Controls:
Species Specificity: human	WB : HepG2 cells, HEK-293 cells, Jurkat cells, HeLa cells
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	IHC : human stomach cancer tissue,

## Background Information

UQCRC2 (Ubiquinol-cytochrome-c reductase complex core protein 2) is also named as complex III subunit 2, core protein II and belongs to the UQCRC2/QCR2 subfamily. The bc1 complex contains 11 subunits: 3 respiratory subunits (cytochrome b, cytochrome c1 and Rieske/UQCRFS1), 2 core proteins (UQCRC1/QCR1 and UQCRC2/QCR2) and 6 low-molecular weight proteins (UQCRH/QCR6, UQCRB/QCR7, UQCRQ/QCR8, UQCR10/QCR9, UQCR11/QCR10 and a cleavage product of Rieske/UQCRFS1) and is part of the mitochondrial respiratory chain. This protein distributes in mitochondrion inner membrane, peripheral membrane protein and matrix side.


## Storage

Storage:  
Store at -20°C. Stable for one year after shipment.  
Storage Buffer:  
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83667-2-RR (UQCRC2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 83667-2-RR (UQCRC2 antibody) at dilution of 1:200 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer(pH9).



1x10<sup>6</sup> HepG2 cells were intracellularly stained with 0.25 ug UQCRC2 Recombinant antibody (83667-2-RR, Clone:240454D9) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (red), or 0.25 ug Isotype Control (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).



Biolayer interferometry (BLI) kinetic assays of 83667-2-RR against Human UQCRC2 were performed. The affinity constant is 21.8 nM.