For Research Use Only

## TNF-alpha Recombinant antibody

Catalog Number:83721-7-RR





Basic Information	Catalog Number: 83721-7-RR	GenBank Accession Number: NM_000594.4	Purification Method: Protein A purfication
	Concentration: 1000 ug/ml	GeneID (NCBI): 7124	CloneNo.: 240668B4
	Source: Rabbit	ENSEMBL Gene ID: ENSG00000232810	Recommended Dilutions: WB 1:2000-1:10000
	lsotype: IgG	UNIPROT ID: P01375	
		Full Name: tumor necrosis factor (TNF superfamily, member 2) Calculated MW: 26 kDa	
		Applications	Tested Applications:
WB, ELISA WB : THP-1 cells, LPS treated THP-1 cells Species Specificity: human			-1 cells, LPS treated THP-1 cells
Background Information	TNF, as also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, ins resistance, and cancer.		
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azid Aliquoting is unnecessary for	e and 50% glycerol pH 7.3.	

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data





Untreated, LPS and Brefeldin A treated THP-1 cells were subjected to SDS PAGE followed by western blot with 83721-7-RR (TNF-alpha antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. Biolayer interferometry (BLL) kinetic assays of 83721-7-RR against Human TNF-alpha were performed. The affinity constant is 8.51 nM.