For Research Use Only

CX3CL1 Recombinant antibody

Catalog Number:83738-5-RR




Basic Information	Catalog Number: 83738-5-RR	GenBank Accession Number: NM_009142.3	Purification Method: Protein A purfication	
	Concentration: 1000 ug/ml	GeneID (NCBI): 20312	CloneNo.: 240738E2	
	Source: Rabbit	UNIPROT ID: O35188	Recommended Dilutions: WB 1:500-1:2000	
	Isotype: IgG	Full Name:IHC 1:500-1:2000chemokine (C-X3-C motif) ligand 1		
		Calculated MW: 42 kDa		
		Observed MW: 55-75 kDa		
Applications	Tested Applications: WB, IHC, ELISA		Positive Controls:	
	Species Specificity:		se brain tissue, mouse lung tissue se brain tissue,	
	Note-IHC: suggested al TE buffer pH 9.0; (*) Al retrieval may be perfo buffer pH 6.0	lternatively, antigen		
Background Information	CX3CL1, also known as fractalkine, is a trans-membrane chemokine that acts as a ligand for both CX3CR1 and integrins $\alpha v \beta 3$ and $\alpha 4 \beta 1$ (PMID: 9479488; 23125415). CX3CL1 is expressed by many hematopoietic and nonhematopoietic cells, including macrophages, endothelial cells, neurons, and fibroblasts. It is involved in immune responses and inflammation (PMID: 21045240). CX3CL1 exists in a membrane-anchored molecule and a soluble form, generated by proteolytic cleavage (PMID: 11495925; 12714508).			
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azio Aliquoting is unnecessary fo	de and 50% glycerol pH 7.3.		

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data





WVW PT CELIES OM

Various lysates were subjected to SDS PAGE followed by western blot with 83738-5-RR (Cx3cl1 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83738-5-RR (CX3CL1 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 83738-5-RR (CX3CL1 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLL) kinetic assays of 83738-5-RR against Mouse CX3CL1 were performed. The affinity constant is 0.744 nM.