

# NRAS Recombinant monoclonal antibody, PBS Only

Catalog Number: 83815-2-PBS

## Basic Information

<b>Catalog Number:</b> 83815-2-PBS	<b>GenBank Accession Number:</b> BC005219	<b>Purification Method:</b> Protein A purification
<b>Source:</b> Rabbit	<b>GeneID (NCBI):</b> 4893	<b>CloneNo.:</b> 242810G11
<b>Isotype:</b> IgG	<b>UNIPROT ID:</b> P01111	
<b>Immunogen Catalog Number:</b> AG1081	<b>Full Name:</b> neuroblastoma RAS viral (v-ras) oncogene homolog	
	<b>Calculated MW:</b> 21 kDa	
	<b>Observed MW:</b> 21 kDa	

## Applications

**Tested Applications:**  
WB, IHC, Indirect ELISA

**Species Specificity:**  
human, mouse, rat

## Background Information

NRAS, also named as N-ras and NRAS1, is neuroblastoma RAS viral (v-ras) oncogene homolog from the mammalian ras gene family and it is a member of the small GTPase superfamily. RAS proteins are involved in signal transduction pathways, and bind GDP/GTP and possess intrinsic GTPase activity. It is mapped on chromosome 1, and it is activated in HL60, a promyelocytic leukemia line. Defects in NRAS are a cause of juvenile myelomonocytic leukemia (JMML). NRAS is one member of RAS gene family of oncoproteins, which is commonly mutated in melanoma and hematopoietic cancers via mapped on chromosome 1 (PMID: 2327491, PMID: 26990546). NRAS mediates activation of both mitogen-activated protein kinase (MAPK) and PI3K/AKT/MYC signaling (PMID: 17297468). NRAS induced classical MAPK signaling leads to cyclin D1 expression and cell cycle dysregulation and promotion of pro-survival pathways (PMID: 7970723, PMID: 18246127). In addition, NRAS effectively prevents Glycogen Synthase Kinase3 (GSK3)-mediated phosphorylation of MYC via PI3K/AKT, which results in enhanced activity of endogenous MYC protein (PMID: 17297468). Mutational NRAS causes Ras-GTP to be in a state of continuous activation, which results in malignant proliferation and metastasis (PMID: 24985059).

## Storage

**Storage:**  
Store at -80°C.

**The product is shipped with ice packs. Upon receipt, store it immediately at -80°C**

**Storage Buffer:**  
PBS only, pH7.3


For technical support and original validation data for this product please contact:

T: 4006900926


E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83815-2-RR (NRAS antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83815-2-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse pancreas tissue slide using 83815-2-RR (NRAS antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83815-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 83815-2-RR (NRAS antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83815-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue slide using 83815-2-RR (NRAS antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83815-2-PBS in a different storage buffer formulation.



Biolayer interferometry (BLI) kinetic assays of 83815-2-RR against Human NRAS were performed. The affinity constant is 0.152 nM.