For Research Use Only

Mouse ASC/TMS1 Recombinant antibody, PBS Only (Capture)

Catalog Number:83858-3-PBS



Purification Method:

Protein A purification

CloneNo.:

240908D5

Basic Information

Catalog Number: GenBank Accession Number: 83858-3-PBS NM_023258

 Concentration:
 GeneID (NCBI):

 1 mg/ml
 66824

 Source:
 UNIPROT ID:

Source: UNIPROT II
Rabbit Q9EPB4
Isotype: Full Name:

IgG PYD and CARD domain containing

Immunogen Catalog Number: Calculated MW:

AG32519 21KD

Observed MW: 21-25 kDa

Applications

Tested Applications:

WB, IHC, Cytometric bead array, Indirect ELISA

Species Specificity:

mouse

Background Information

PYCARD (PYD and CARD domain containing), also known as ASC and CARD5. The calculated molecular weight of PYCARD is 21 kDa. It is located in cytosol, extracellular region and nucleus. PYCARD is also expressed in several structures, including alimentary system; brain; heart; hemolymphoid system gland; and urinary system. The gene enables identical protein binding activity and protein dimerization activity. Involved in several processes, including activation of cysteine-type endopeptidase activity; positive regulation of cytokine production; and regulation of defense response. It acts upstream of or within several processes, including defense response to Grampositive bacterium; positive regulation of macromolecule metabolic process; and regulation of autophagy. It is also a part of inflammasome complex. Finally, it orthologous to human PYCARD (PYD and CARD domain containing).

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83858-3-RR (PYCARD antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83858-3-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 83858-3-RR (ASC/TMS1 antibody) at dilution of 1:500 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83858-3-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded mouse colon tissue slide using 83858-3-RR (ASC/TMS1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83858-3-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP00782-1, MOUSE ASC/TMS1 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 83858-3-PBS. Detection antibody: 83858-1-PBS. Standard: Ag32519. Range: 0.47-60 ng/mL



Cytometric bead array standard curve of MP00782-2, MOUSE ASC/TMS1 Recombinant Matched Antibody Pair, PBS Only. Capture antibody: 83858-3-PBS. Detection antibody: 83858-4-PBS. Standard: Ag32519. Range: 0.47-60 ng/mL



Biolayer interferometry (BLI) kinetic assays of 83858-3-RR against Mouse PYCARD were performed. The affinity constant is 3.58 nM.