For Research Use Only

Loricrin Recombinant antibody

Catalog Number:83865-1-RR



Basic Information

Catalog Number: GenBank Accession Number: 83865-1-RR NM_000427

Concentration: GeneID (NCBI): 1000 ug/ml 4014

Source: UNIPROT ID: Rabbit P23490

Isotype: Full Name: IgG loricrin

Calculated MW: 26 kDa

Purification Method: Protein A purfication

CloneNo.:

240605C8

Recommended Dilutions: IHC 1:250-1:1000 IF/ICC 1:125-1:500

Applications

Tested Applications: IHC, IF/ICC, FC (Intra), ELISA

Species Specificity: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

IHC: mouse skin tissue,
IF/ICC: A431 cells,

Background Information

Loricrin, also named LOR and LRN, is a major component of the cornified cell envelope (CE) expressed late in epidermal differentiation in the granular layers of normal skin. This antibody is specific to Loricrin.

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed A431 cells using LOR antibody (83865-1-RR, Clone: 240605C8) at dilution of 1:250 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



1x10^6 A431 cells were intracellularly stained with 0.25 ug Loricrin Recombinant antibody (83865-1-RR, Clone:240605C8) and APC-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)(red), or 0.25 ug Isotype Control (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using 83865-1-RR (Loricrin antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse skin tissue slide using 83865-1-RR (Loricrin antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).