

For Research Use Only

NCAM1/CD56 Recombinant antibody

Catalog Number: 83883-2-RR



Basic Information

Catalog Number: 83883-2-RR	GenBank Accession Number: GeneID (NCBI): 17967	Purification Method: Protein A purification
Concentration: 1000 ug/ml	UNIPROT ID: P13595-1	CloneNo.: 240936B9
Source: Rabbit	Full Name: neural cell adhesion molecule 1	Recommended Dilutions: WB 1:5000-1:50000
Isotype: IgG	Calculated MW: 119 kDa	
	Observed MW: 120 kDa, 140 kDa, 180 kDa	

Applications

Tested Applications: WB, ELISA	Positive Controls:
Species Specificity: mouse, rat	WB : mouse cerebellum tissue, C6 cells, mouse brain tissue, rat brain tissue

Background Information

Neural cell adhesion molecule 1 (NCAM1, also known as CD56) is a cell adhesion glycoprotein of the immunoglobulin (Ig) superfamily. It is a multifunction protein involved in synaptic plasticity, neurodevelopment, and neurogenesis. NCAM1 is expressed on human neurons, glial cells, skeletal muscle cells, NK cells, and a subset of T cells, and the expression is observed in a wide variety of human tumors, including myeloma, myeloid leukemia, neuroendocrine tumors, Wilms' tumor, neuroblastoma, and NK/T cell lymphomas. Three major isoforms of NCAM1, with molecular masses of 120, 140, and 180 kDa, are generated by alternative splicing of mRNA (PMID: 9696812). The glycosylphosphatidylinositol (GPI)-anchored NCAM120 and the transmembrane NCAM140 and NCAM180 consist of five Ig-like domains and two fibronectin-type III repeats (FNIII). All three forms can be posttranslationally modified by the addition of polysialic acid (PSA) (PMID: 14976519). Several other isoforms have also been described (PMID: 1856291).

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 83883-2-RR (NCAM1/CD56 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Biolayer interferometry (BLI) kinetic assays of 83883-2-RR against Mouse Ncam1 were performed. The affinity constant is 0.833 nM.