

For Research Use Only

FBP2 Recombinant antibody, PBS Only

Catalog Number: 83910-5-PBS



Basic Information

Catalog Number:

83910-5-PBS

Size:

1 mg/ml

Source:

Rabbit

Isotype:

IgG

Immunogen Catalog Number:

AG18245

GenBank Accession Number:

BC113632

GeneID (NCBI):

8789

UNIPROT ID:

O00757

Full Name:

fructose-1,6-bisphosphatase 2

Calculated MW:

339 aa, 37 kDa

Observed MW:

37 kDa

Purification Method:

Protein A purification

CloneNo.:

240879C9

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Species Specificity:

human, mouse, rat

Background Information

FBP2, also named muscle FBP, belongs to the FBPase class 1 family. FBP2 is a ubiquitously expressed enzyme of glycogen synthesis from non-carbohydrates, e.g., lactate (glyconeogenesis). However, it also plays a variety of non-enzymatic functions. FBP2 is involved in the regulation of hypoxia-inducible factor 1 (Hif1) stability, cell cycle-dependent events, biogenesis of mitochondria, and protection of the organelles against high reactive oxygen species (ROS)- and high Ca²⁺-induced stress (PMID: 32962293, PMID: 35626746). The calculated molecular weight of FBP2 is 37 kDa.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Mouse heart tissue were subjected to SDS PAGE followed by western blot with 83910-5-RR (FBP2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 83910-5-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 83910-5-RR (FBP2 antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83910-5-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 83910-5-RR (FBP2 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83910-5-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human ovary cancer tissue slide using 83910-5-RR (FBP2 antibody) at dilution of 1:600 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 83910-5-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse skeletal muscle tissue using FBP2 antibody (83910-5-RR, Clone: 240879C9) at dilution of 1:250 and CoraLite®488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2). This data was developed using the same antibody clone with 83910-5-PBS in a different storage buffer formulation.



Bi-layer interferometry (BLI) kinetic assays of 83910-5-RR against Human FBP2 were performed. The affinity constant is 4.67 nM.