For Research Use Only

transgelin/SM22 Recombinant antibody

Catalog Number:83922-2-RR



Basic Information

Catalog Number: GenBank Accession Number: 83922-2-RR BC004927

 Concentration:
 GeneID (NCBI):
 CloneNo.:

 1000 ug/ml
 6876
 240864A11

 Source:
 UNIPROT ID:
 Recommended Dilutions:

 Rabbit
 Q01995
 WB 1:5000-1:50000

 Isotype:
 Full Name:
 IHC 1:50-1:500

 IgG
 transgelin
 IF-P 1:200-1:800

AG0764 22 kDa

Observed MW: 19-22 kDa

Calculated MW:

Applications

Tested Applications: WB, IHC, IF-P, ELISA Species Specificity: human, mouse, rat

Immunogen Catalog Number:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Positive Controls:

WB: mouse colon tissue, rat colon tissue, mouse

Purification Method:

Protein A purfication

stomach tissue, rat stomach tissue

IHC: mouse heart tissue, IF-P: mouse heart tissue,

Background Information

The transgelin family is a protein group belonging to the 22kd actin-related calponin superfamily. Of all three isoforms, transgelin 1 is the best characterized. Transgelin 1, or SM22alpha, is a specific marker for differentiated smooth muscle cells. Transgelin 2, also known as SM22 beta, is expressed by both smooth and non-smooth muscle cells in a temporally and spatially regulated pattern. Trangenlin 3, also known as NP25, is only found in highly differentiated neuronal cells. This antibody was generated against full-length transgelin 1 protein. It can cross-react with other two transgenes based on the sequence similarity.

Storage


Storage

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 83922-2-RR (transgelin/SM22 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 83922-2-RR (transgelin/SM22 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 83922-2-RR (transgelin/SM22 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse heart tissue using transgelin/SM22 antibody (83922-2-RR, Clone: 240864A11) at dilution of 1:400 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse heart tissue using transgelin/SM22 antibody (83922-2-RR, Clone: 240864A11) at dilution of 1:400 and Coralite@488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 83922-2-RR against Human transgelin/SM22 were performed. The affinity constant is 14.2 nM.