For Research Use Only

SP100 Recombinant antibody

Catalog Number:84012-5-RR




| Basic Information | Catalog Number: 84012-5-RR | GenBank Accession Number BC011562 | r: Purification Method: Protein A purfication | |
|------------------------|--|--|--|--|
| | Concentration: 1000 µg/ml | GeneID (NCBI): 6672 | CloneNo.: 241153F8 | |
| | Source: Rabbit Isotype: IgG Immunogen Catalog Number: AG35193 | UNIPROT ID: P23497 Full Name: SP100 nuclear antigen | Recommended Dilutions: WB 1:2000-1:10000 | |
| | | | IHC 1:400-1:1600 IF/ICC 1:125-1:500 | |
| | | Calculated MW: 879 aa, 100 kDa | | |
| | | Observed MW: 70-100 kDa, 110-140 kDa | | |
| Applications | Tested Applications: | Posi | tive Controls: | |
| | WB, IHC, IF/ICC, ELISA Species Specificity: human | | WB: A431 cells, HeLa cells | |
| | | | : human ovary cancer tissue, | |
| | Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 | | | |
| Background Information | SP100 nuclear antigen, belongs to the nuclear bodies(NBs), has a role in the control of gene expression. It contains a HSR domain, which is important for the nuclear body targeting as well as for the dimerization, and one PxVxL motif, which if required for interaction with chromoshadow domains. Sp100 is an acidic protein that exerts a highly aberrant electrophoretic mobility in SDS-polyacrylamide gel electrophoresis. | | | |
| Storage | Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.02% sodium azide and Aliquoting is unnecessary for -20° | i 50% glycerol, pH7.3 | | |


For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84012-5-RR (SP100 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 84012-5-RR (SP100 antibody) at dilution of 1:800 (under 10x lens). Heat mediated antigen retrieval with.



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 84012-5-RR (SP100 antibody) at dilution of 1:800 (under 40x lens). Heat mediated antigen retrieval with.



Immunofluorescent analysis of (4% PFA) fixed A431 cells using SP100 antibody (84012-5-R, Clone: 241153F8) at dilution of 1:250 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed A431 cells using SP100 antibody (84012-5-RR, Clone: 241153F8) at dilution of 1:250 and CoraLite@488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red).