

For Research Use Only

CD98/SLC3A2 Recombinant monoclonal antibody

Catalog Number: 84040-7-RR



Basic Information

Catalog Number:	84040-7-RR	GenBank Accession Number:	NM_008577.4	Purification Method:	Protein A purification
Concentration:	1000 µg/ml	GenID (NCBI):	17254	CloneNo.:	241247F12
Source:	Rabbit	UNIPROT ID:	P10852-1	Recommended Dilutions:	WB: 1:5000-1:50000 IHC: 1:1000-1:4000
Isotype:	IgG	Full Name:	solute carrier family 3 (activators of dibasic and neutral amino acid transport), member 2		
Immunogen Catalog Number:	EG1374	Calculated MW:	58 kDa		

Applications

Tested Applications:	WB, IHC, ELISA	Positive Controls:	
Species Specificity:	mouse	WB:	mouse kidney tissue, mouse testis tissue
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			IHC: mouse brain tissue, mouse kidney tissue

Background Information

CD98 is a cell-surface heterodimer consisting of a heavy chain (CD98hc) and a light chain. CD98hc also interacts with certain integrin β -subunits to regulate cell migration, survival, proliferation, and adhesion/polarity. CD98hc is overexpressed on the cell surface of many cancers and increased CD98hc expression is associated with the development and progression of tumors. (PMID:25084765). This protein has 4 isoforms with the molecular mass of 58-71 kDa and can be detected 85-94 kDa due to glycosylation. The glycosylated CD98hc can link to a non-glycosylated light chain (~40 kDa) via a disulfide bond to form a heterodimeric CD98 antigen with molecular mass of 120-130 kDa (PMID: 14770309).

Storage


Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol, pH7.3
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




mouse kidney tissue were subjected to SDS PAGE followed by western blot with 84040-7-RR (CD98 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




mouse testis tissue were subjected to SDS PAGE followed by western blot with 84040-7-RR (CD98 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 84040-7-RR (CD98 antibody) at dilution of 1:2000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 84040-7-RR (CD98 antibody) at dilution of 1:2000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Biolayer interferometry (BLI) kinetic assays of 84040-7-RR against Mouse CD98/SLC3A2 were performed. The affinity constant is 1.36 nM.