For Research Use Only

CD38 Recombinant antibody

Catalog Number:84070-2-RR




Basic Information	Catalog Number: 84070-2-RR	GenBank Accession Number: BC007964	Purification Method: Protein A purification				
	Size: 1000 ug/ml	GeneID (NCBI): 952	CloneNo.: 241246B1				
	Source: Rabbit	ENSEMBL Gene ID: ENSG0000004468	Recommended Dilutions: WB 1:5000-1:50000				
	Isotype: IgG	UNIPROT ID: P28907	IF/ICC 1:125-1:500				
		Full Name: CD38 molecule Calculated MW: 300 aa, 34 kDa Observed MW: 42 kDa					
				Applications	Tested Applications:	Positive	Controls:
					WB, IF/ICC, ELISA Species Specificity: human	WB : Raji	WB : Raji cells, A549 cells, MOLT-4 cells, Ramos cells
IF/ICC : I	Raji cells,						
Background Information	CD38, also known as ADP-ribosyl cyclase 1, is a type II transmembrane glycoprotein with a short N-terminal cytoplasmic tail, a single membrane-spanning domain, and a C-terminal extracellular region with four N-glycosylation sites (PMID: 2319135). The extracellular domain of CD38 has bifunctional enzyme activities that catalyze synthesis of cyclic ADP ribose from nicotinamide adenine dinucleotide (NAD) and hydrolysis of cyclic ADP ribose (PMID: 10636863). CD38 is expressed on a variety of hematopoietic and non-hematopoietic cells and is involved in diverse processes such as generation of calcium-mobilizing metabolites, cell activation, and chemotaxis (PMID: 25938500).						
Storage	Storage: Store at -20°C. Stable for one Storage Buffer: PBS with 0.02% sodium azide Aliquoting is unnecessary for	e and 50% glycerol pH 7.3.					


For technical support and original validation data for this product please contact: T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 84070-2-RR (CD38 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WWW.ETGLAE.COM

Immunofluorescent analysis of (4% PFA) fixed Raji cells using CD38 antibody (84070-2-RR, Clone: 241246B1) at dilution of 1:250 and CoraLite®488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). Immunofluorescent analysis of (4% PFA) fixed Raji cells using CD38 antibody (84070-2-RR, Clone: 241246B1) at dilution of 1:500 and CoraLite®594-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-4).



Biolayer interferometry (BLl) kinetic assays of 84070-2-RR against Human CD38 were performed. The affinity constant is 0.999 nM.