## For Research Use Only

## EPCAM/CD326 Recombinant antibody

Catalog Number:84073-6-RR 1 Publications



**Basic Information** 

Catalog Number: 84073-6-RR Concentration: 1000 ug/ml

Source: Rabbit Isotype:

Immunogen Catalog Number:

EG1370

GenBank Accession Number:

NM\_002354.3 GeneID (NCBI):

**UNIPROT ID:** P16422 Full Name:

35-40 kDa

4072

epithelial cell adhesion molecule

Calculated MW: 35 kDa Observed MW:

Positive Controls:

WB: A431 cells, HCT 116 cells, HaCaT cells, MCF-7

**Purification Method:** 

Protein A purfication

WB 1:5000-1:50000 IHC 1:500-1:2000

IF/ICC 1:200-1:800

Recommended Dilutions:

CloneNo.:

241243B10

IHC: human ovary cancer tissue,

IF/ICC: MCF-7 cells,

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA Cited Applications:

IF

Species Specificity:

human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

**Background Information** 

Epithelial cell adhesion molecule (EpCAM, CD326) is a type I transmembrane glycoprotein that functions as a homophilic, epithelial-specific intercellular cell-adhesion molecule. In addition to cell adhesion, EpCAM is also involved in cellular signaling, cell migration, proliferation, and differentiation. EpCAM is highly expressed on most carcinomas and therefore of potential use as a diagnostic and prognostic marker for a variety of carcinomas, and has become a therapeutic target. EpCAM may occur in distinct forms due to glycosylation. (PMID: 20837599; 19249674; 21576002; 22647938; 12691820)

**Notable Publications** 

Author **Pubmed ID** Journal Application Wenzhao Su 39558398 J Transl Med

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 84073-6-RR (EPCAM/CD326 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 84073-6-RR (EPCAM/CD326 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human ovary cancer tissue slide using 84073-6-RR (EPCAM/CD326 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using EPCAM/CD326 antibody (84073-6-RR, Clone: 241243B10 ) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2).



Biolayer interferometry (BLL) kinetic assays of 84073-6-RR against Human EPCAM/CD326 were performed. The affinity constant is 2.56 nM.