For Research Use Only

CDC25A Recombinant antibody, PBS Only

Catalog Number:84145-2-PBS



Purification Method:

Protein A purfication

CloneNo.:

240813F3

Basic Information

Catalog Number: 84145-2-PBS

Concentration: 1 mg/ml

Source: UNIPROT ID: Rabbit P30304
Isotype: Full Name:

G cell division cycle 25 homolog A (S.

pombe)

NM_001789

GeneID (NCBI):

Calculated MW: 59 kDa

GenBank Accession Number:

Applications

Tested Applications: IHC, IF/ICC, ELISA Species Specificity: human

Background Information

CDC25A belongs to the MPI phosphatase family. It is a tyrosine protein phosphatase which functions as a dosage-dependent inducer of mitotic progression. CDC25A directly dephosphorylates CDC2 and stimulates its kinase activity. It also dephosphorylates CDK2 in complex with cyclin E, in vitro. CDC25A overexpression in human cancers might contribute to tumorigenesis. This protein can be phosphorylated to reveal the molecular weight from 55 kDa to 70 kDa(PMID:21376736; 8156993). The antibody is specific to CDC25A.

Storage

Storage: Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CDC25A antibody (84145-2-RR, Clone: 240813F3) at dilution of 1:250 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Rabbit 1gG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84145-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 84145-2-RR (CDC25A antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84145-2-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human stomach cancer tissue slide using 84145-2-RR (CDC25A antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 84145-2-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using CDC25A antibody (84145-2-RR, Clone: 240813F3) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Rabbit IgG(H+L) (SA00013-2), CL594-Phalloidin (red). This data was developed using the same antibody clone with 84145-2-PBS in a different storage buffer formulation.